

EXPRESS MAIL LABEL NO: \_\_\_\_\_  
Docket No. P-2762-US1

**AN ISOLATED NUCLEIC ACID ENCODING P-HYDE PROTEIN  
AND METHODS OF INDUCING SUSCEPTIBILITY  
TO INDUCTION OF CELL DEATH IN CANCER**

5 This application is a continuation-in-part of U.S. Serial No. 09/302,457, filed April 29, 1999, the contents of which are hereby incorporated by reference.

**FIELD OF INVENTION**

10 This invention provides an isolated nucleic acid of the p-Hyde gene, analogs, fragments, mimetics, mutants, synthetics, and variants thereof. This invention is directed to a method of inducing susceptibility to apoptosis with p-Hyde, a method of suppressing tumor growth with p-Hyde, and a method of treating a subject with cancer with p-Hyde alone or in combination with radiation, chemotherapy, or UV mimetic drugs.

15 **BACKGROUND OF THE INVENTION**

Prostate cancer is the most common malignancy in men with over 317,000 new cases and the second leading cause of male cancer deaths in the United States (Boring et al., 1993; Steiner et. al, 1995). The molecular mechanisms responsible for the development, progression, and metastasis of prostate cancer remain largely unknown. Up to 20% of prostate cancers occur in men under the age of 65 years of age (Silverberg, 1986) suggesting that prostate carcinogenesis is not only associated with aging, but also to hereditary factors (Silverberg, 1987; McLellan and Norman, 1995; Carter et al., 1992). Genetic linkage studies of 691 affected families have revealed that an earlier age of onset of the disease in the proband and the presence of multiple affected family members are important determinants that increase the risk of prostate cancer. The pattern of inheritance of the putative prostate cancer gene appears to be autosomal dominant with an 88% penetrance rate (Steinberg, 1990). Thus, hereditary factors play an important role in prostate oncogenesis.

30 Like many carcinomas, prostate cancer formation is a multistep process involving tumor initiation, promotion, conversion, and progression (Carter et al., 1990; Sandberg, 1992).

5 This process is driven by chromosomal instability, spontaneous mutations, and carcinogen  
induced genetic and epigenetic changes. *Chromosomal instability* leads to the total or partial  
gain or loss of chromosomes, translocations, and other abnormalities. *Spontaneous*  
10 *mechanisms* are age-related and include activation of oncogenes or inactivation of tumor  
suppressor genes by genetic mutations. These mutations result in the misincorporation of  
nucleotides during DNA replication of the coding region, alteration of the intron-exon  
junction sequences affecting the splicing mechanism, and aberrations of regulatory sequences  
changing the control of critical genes. These mutations escape genetic surveillance by a  
battery of DNA repair mechanisms and its associated gene products, such as p53 (Effert et  
al., 1992, Isaacs et al, 1991; Mellon et al., 1992) and p21 (El-Deiry et al., 1994) and PCNA  
15 (Templeton et al., 1996). *Carcinogen-induced genetic and epigenetic changes* initiate  
tumors as a consequence of the direct damaging effects of carcinogenic agents of the DNA  
altering gene expression. Tumor initiation is subsequently followed by tumor promotion as  
affected cells have selective reproductive and clonal expansion capabilities through altered  
signaling transduction and proliferation responses to growth factors, resistance to  
20 cytotoxicity, and deregulation of terminal differentiation (Yuspa and Poirier, 1988;  
Weinstein, 1987). Finally, tumor promotion is succeeded by other genetic mutational events  
that lead to loss of hormone sensitivity, increased cell motility, invasion, alterations in  
programmed cell death and metastasis. Accordingly, the initiation and progression of cancer  
is a multistep process whereby genetic alterations or mutations of critical genes ultimately  
25 dictate defined cell phenotypes which differ in regard to many important cellular activities  
including cell proliferation, differentiation, and programmed cell death. The exact mutational  
events responsible for the multistep progression of prostate cancer, however, is unknown.  
A better understanding of the molecular mechanisms responsible for prostate cancer may  
lead to new therapies to combat, and perhaps, to even prevent prostate cancer.

30

#### SUMMARY OF THE INVENTION

The present invention provides an isolated nucleic acid encoding a P-Hyde gene. The pHyde  
gene as shown herein is associated with: (1) the regression of tumor growth *in vivo* (2) the  
induction to susceptibility to apoptosis caused by UV or chemotherapy induced DNA

5 damage, and (3) prevention of DNA repair with the upregulation of apoptosis as the result of UV damage and the failure to repair DNA.

10 This invention provides a novel class of genes which act as inhibitor of a DNA repair enzyme and induce susceptibility of cancer cells to cell death. Also, this invention provides an isolated nucleic acid which encodes a mammalian p-Hyde protein which induces susceptibility of a cancer cell to cell death, including allelic, analogs, fragments, mimetics, mutants, synthetics, or variants thereof. This invention provides an isolated nucleic acid which encodes a human p-Hyde protein which induces susceptibility of a cancer cell to cell death, including allelic, analogs, fragments, mimetics, mutants, synthetics, or variants thereof.

15 This invention provides a vector comprising the isolated nucleic acid encoding P-Hyde gene. This invention provides a replication-defective recombinant E1/E3 deleted adenovirus containing a truncated RSV promoter and the P-Hyde cDNA gene (AdRSVpHyde).

20 This invention provides an oligonucleotide of at least 15 nucleotides capable of specifically hybridizing with a sequence of nucleotides present within a nucleic acid which encodes the human p-Hyde, or a sequence which is complementary to the nucleic acid which encodes the human p-Hyde. This invention provides an antisense molecule, triplex oligonucleotide, or ribozyme which is capable of specifically hybridizing with the isolated nucleic acid encoding p-Hyde.

25 This invention provides a method for producing a polypeptide which comprises growing the host vector system under suitable conditions permitting production of the polypeptide and recovering the polypeptide so produced. In one embodiment the method of obtaining a polypeptide in purified form comprises: (a) introducing the vector into a suitable host cell; (b) culturing the resulting cell so as to produce the polypeptide; (c) recovering the polypeptide produced in step (b); and (d) purifying the polypeptide so recovered.

30 This invention provides a polypeptide comprising the amino acid sequence of a human p-

5 Hyde. This invention provides a fusion protein or chimeric comprising the polypeptide. This invention provides an antibody which specifically binds to the polypeptide. This invention provides a pharmaceutical composition comprising an amount of the polypeptide and a pharmaceutically effective carrier or diluent.

10 This invention provides a method for determining whether a subject carries a mutation in the p-Hyde gene which comprises: (a) obtaining an appropriate nucleic acid sample from the subject; and (b) determining whether the nucleic acid sample from step (a) is, or is derived from, a nucleic acid which encodes mutant p-Hyde so as to thereby determine whether a subject carries a mutation in the p-Hyde gene. In one embodiment the nucleic acid sample  
15 in step (a) comprises mRNA corresponding to the transcript of DNA encoding a mutant p-Hyde, and wherein the determining of step (b) comprises: (i) contacting the mRNA with the oligonucleotide under conditions permitting binding of the mRNA to the oligonucleotide so as to form a complex; (ii) isolating the complex so formed; and (iii) identifying the mRNA in the isolated complex so as to thereby determine whether the mRNA is, or is derived from,  
20 a nucleic acid which encodes mutant p-Hyde.

This invention provides a method for screening a tumor sample from a human subject for a somatic alteration in a p-Hyde gene in said tumor which comprises gene comparing a first sequence selected from the group consisting of a p-Hyde gene from said tumor sample, p-Hyde RNA from said tumor sample and p-Hyde cDNA made from mRNA from said tumor sample with a second sequence selected from the group consisting of p-Hyde gene from a nontumor sample of said subject, p-Hyde RNA from said nontumor sample and p-Hyde cDNA made from mRNA from said nontumor sample, wherein a difference in the sequence of the p-Hyde gene, p-Hyde RNA or p-Hyde cDNA from said tumor sample from the  
25 sequence of the p-Hyde gene, p-Hyde RNA or p-Hyde cDNA from said nontumor sample indicates a somatic alteration in the p-Hyde gene in said tumor sample.  
30

This invention provides a method for screening a tumor sample from a human subject for the presence of a somatic alteration in a p-Hyde gene in said tumor which comprises comparing  
35 p-Hyde polypeptide from said tumor sample from said subject to p-Hyde polypeptide from



5 a nontumor sample from said subject to analyze for a difference between the polypeptides,  
wherein said comparing is performed by (i) detecting either a full length polypeptide or a  
truncated polypeptide in each sample or (ii) contacting an antibody which specifically binds  
to either an epitope of an altered p-Hyde polypeptide or an epitope of a wild-type p-Hyde  
polypeptide to the p-Hyde polypeptide from each sample and detecting antibody binding,  
10 wherein a difference between the p-Hyde polypeptide from said tumor sample from the p-  
Hyde polypeptide from said nontumor sample indicates the presence of a somatic alteration  
in the p-Hyde gene in said tumor sample.

This invention provides a method for identifying a chemical compound which is capable  
15 inducing susceptibility to cell death which comprises: (a) contacting the p-Hyde with a  
chemical compound under conditions permitting binding between the p-Hyde and the  
chemical compound; (b) detecting specific binding of the chemical compound to the p-Hyde;  
and (c) determining whether the chemical compound inhibits the p-Hyde so as to identify a  
chemical compound which is capable of capable inducing susceptibility to cell death.

20 This invention provides a method of inhibiting the growth of cancer cells, comprising the  
steps of obtaining the cells and contacting the cells of the subject with a replication-deficient  
adenovirus type 5 expression vector comprising an adenovirus genome having a deletion in  
the E1 and E3 region of the genome and an insertion within the region of a nucleic acid  
25 encoding p-Hyde under the control of a Rous Sarcoma virus promoter, thereby inhibiting the  
growth of the prostate cancer cells.

This invention provides a method of inhibiting the growth of prostate cancer cells,  
comprising: 1) obtaining a sample of prostate cells from a subject; 2) contacting the cells  
30 with a replication deficient adenovirus type 5 expression vector which comprises an  
adenovirus genome having a deletion in the E1 and E3 regions of the genome and an  
insertion within the regions of a p-Hyde cDNA under the control of a Rous Sarcoma virus  
promoter; and 3) introducing the cells into the subject, thereby inhibiting the growth of the  
cancer cells.

35

5 This invention provides a method of suppressing the growth of cancer cells in a subject, comprising introducing into the cancer cell an amount of a nucleic acid encoding a p-Hyde protein, a nucleic acid encoding a fragment of p-Hyde protein, or the nucleic acid encoding a mutant p-Hyde protein, thereby suppressing the growth of cancer cells in the subject.

10 This invention provides a method of suppressing growth of cancer cells in a subject, comprising administering to the subject a pharmaceutical composition comprising a therapeutically effective amount of a nucleic acid encoding a p-Hyde protein, a nucleic acid encoding a fragment of p-Hyde protein, or the nucleic acid encoding a mutant p-Hyde protein and a pharmaceutical acceptable carrier or diluent, thereby suppressing the growth  
15 of cancer cells in the subject.

This invention provides a method of inducing susceptibility to apoptosis of cancer cells in a subject, comprising introducing into the cancer cell an amount of a nucleic acid encoding a p-Hyde protein, a nucleic acid encoding a fragment of p-Hyde protein, or the nucleic acid  
20 encoding a mutant p-Hyde protein, thereby inducing susceptibility to apoptosis.

This invention provides a method of inducing susceptibility to apoptosis of cancer cells in a subject, comprising administering to the subject a pharmaceutical composition comprising a therapeutically effective amount of a nucleic acid encoding a p-Hyde protein, a nucleic  
25 acid encoding a fragment of p-Hyde protein, or the nucleic acid encoding a mutant p-Hyde protein and a pharmaceutical acceptable carrier or diluent, thereby inducing susceptibility to apoptosis.

This invention provides a method of treating a subject with cancer which comprises  
30 administering to the subject a pharmaceutical composition comprising a therapeutically effective amount of a nucleic acid encoding a p-Hyde protein, a nucleic acid encoding a fragment of p-Hyde protein, or the nucleic acid encoding a mutant p-Hyde protein and a pharmaceutical acceptable carrier or diluent, thereby treating the subject with cancer.

35 This invention provides a method of treating a subject with cancer, comprising: 1)

5 administering to the subject a pharmaceutical composition comprising a therapeutically effective amount of a nucleic acid encoding a p-Hyde protein, a nucleic acid encoding a fragment of p-Hyde protein, or the nucleic acid encoding a mutant p-Hyde protein in combination with radiation, chemotherapy, or UV mimetic drugs; and 2) a pharmaceutical acceptable carrier or diluent, thereby treating the subject with cancer.

10

This invention a method of treating a subject with cancer, comprising administering to the subject a therapeutically effective amount of a pharmaceutical composition comprising: 1) an adenovirus type 5 expression vector which comprises an adenovirus genome having a deletion in the E1 and E3 regions of the genome and an insertion within the regions of a full length sense p-Hyde cDNA under the control of a Rous Sarcoma virus promoter, and 2) a suitable carrier or diluent, thereby treating the subject with cancer. In one embodiment the cancer is selected from a group consisting of: melanoma; lymphoma; leukemia; and prostate, colorectal, pancreatic, breast, brain, or gastric carcinoma.

15

20 Lastly, the present invention provides the means necessary for production of gene-based therapies directed at cancer cells. These therapeutic agents may take the form of polynucleotides comprising all or a portion of the p-Hyde locus placed in appropriate vectors or delivered to target cells in more direct ways such that the function of the p-Hyde protein is reconstituted. Therapeutic agents may also take the form of polypeptides based on either a portion of, or the entire protein sequence of p-Hyde. These may functionally replace the activity of p-Hyde in vivo.

25

#### **BRIEF DESCRIPTION OF THE DRAWINGS**

30

**Figure 1. Schematic presentation of AdRSVpHyde structure.** The 2664 bp inserted fragment contains a 1467 bp full-length pHyde cDNA gene (SEQ ID N0: 1) and 1166 bp 3' untranslated downstream region. The complete sequence of AdRSVpHyde is set forth in Figure 10, Specifically, the nucleic acid sequence of region A in Figure 1 is set forth in

5 Figure 10 Region A and the nucleic acid sequence of region B in Figure 1 is set forth in Figure 10 at Region B.

10 **Figure 2. Expression of pHyde by AdRSVpHyde.** DU145 cells transduced by AdRSVpHyde at MOI=200 were harvested in 48 h post infection for either mRNA or protein extraction. (A) Expression of pHyde at mRNA level in DU145 cells. Sample wells were each loaded with 10 mg of total RNA, electrophoresed in 12.5% agarose gel, transferred to nylon membrane, and hybridized with <sup>32</sup>P-labeled pHyde cDNA. Northern blot was stripped and rehybridized with GAPDH to assess gel loading. (B) Expression of pHyde at protein level in DU145 cells. Protein extracts (50 mg) were loaded on a 12% SDS-PAGE gel. 15 Rabbit anti-rat pHyde antibody was used as the primary antibody.

20 **Figure 3. Inhibitory effects of pHyde on prostate cancer cell growth.** DU145 (A) and LNCaP (B) cells were transduced with or without adenoviral vectors at MOI=100. Cell numbers were counted at day 5 after viral transduction. The data represent the results from two independent experiments with each in duplicates.

25 **Figure 4. AdRSVpHyde inhibits prostate tumor growth *in vivo*.** DU145 cells ( $1.4 \times 10^7$  cells) were injected subcutaneously into the flanks of nude mice. When tumors reached an average volume of 80 mm<sup>3</sup> (about one month after tumor cell inoculation), tumors were untreated (control), or intratumorally injected (day 0) either by  $5 \times 10^9$  pfu control virus AdRSVlacZ (control virus), or  $5 \times 10^9$  pfu AdRSVpHyde (AdRSVpHyde). The tumor sizes were periodically measured at times shown in the figure up to day 52 days post viral injection.

30 **Figure 5. Morphological changes of DU145 and LNCaP cells transduced by AdRSVpHyde.** Cells were transduced by control adenovirus AdRSVlacZ or by AdRSVpHyde at MOI=100. The morphologic features of untreated control cells and viral-transduced cells were recorded at day 5 post viral transduction. All the photos are at the same magnification (66X). (A) and (D): Untreated control cells; (B) and (E): Viral control AdRSVlacZ treated cells; (C) and (F): AdRSVpHyde treated cells. 35

5      **Figure 6. Morphological changes of PC-3, TSU, and PPC-1 cells transduced by**  
AdRSVpHyde. Cells were transduced by control adenovirus AdRSVlacZ or by  
AdRSVpHyde at MOI=100. The morphologic features of untreated control cells and viral-  
transduced cells were recorded at day 5 post viral transduction. All the photos are at the  
same magnification (66X). (A, D, G): Untreated control cells; (B, E, H): Viral control  
10      AdRSVlacZ treated cells; (C, F, I): AdRSVpHyde treated cells.

**Figure 7. Expression of p53 and Rb mRNA in various prostate cancer cell lines.**  
Sample wells were each loaded with 10 mg of total RNA, electrophoresed in 12.5% agarose  
gel, transferred to nylon membrane, and hybridized with <sup>32</sup>P-labeled p53 or Rb cDNA probe.  
15      Northern blot was stripped and rehybridized with GAPDH to normalize the gel loading.

**Figure 8. AdRSVpHyde induced p53 expression in DU145 cells.** The same Northern blot  
in Fig. 2A was stripped and rehybridized with <sup>32</sup>P-labeled p53 cDNA.

20      **Figure 9. AdRSVpHyde induced apoptosis in LNCaP cells.** Cells were untreated or  
transduced by AdRSVpHyde at MOI=100, supernatant were collected 48 h post transduction.  
Soluble DNA was extracted from floating cells and electrophoresed on a 2% agarose gel.

25      **Figure 10. The complete sequence of AdRSVpHyde.**

#### **DETAILED DESCRIPTION OF THE INVENTION**

This invention provides a novel class of genes which act as inhibitor of a DNA repair enzyme  
and induces susceptibility of cancer cells to cell death. Functionally, *P-Hyde* is and  
30      associated with suppression of tumor growth *in vivo* and increased susceptibility to apoptosis  
induced by UV irradiation or FUrD treatment. The upregulation of apoptosis due to UV  
damage is correlated with the presence of intact photoproduct in prostate cancer cell lines  
stably transfected with *p-Hyde*. Use of *p-Hyde* in human gene therapy as monotherapy or in  
combination with radiation or chemotherapy is useful against cancer or hyperproliferative  
35      human diseases. In one embodiment, the class of proteins, such as P-Hyde, are DNA repair



5 enzyme inhibitors which downregulate Nucleotide-Excision-Repair (NER) pathway in prostate cancer, MGMT CNA repair pathway in colon cancer cell line, O<sub>6</sub> methyl guanine methyl transferase enzyme (O<sub>6</sub>MgMT) and 6,4, photoproducts (6,4PP). The class of genes is characterized by comprising a leucine zipper binding domain and a death domain which causes the cell to be apoptotic.

10

This invention provides an isolated nucleic acid which encodes a mammalian p-Hyde protein which induces susceptibility of a cancer cell to cell death, including allelic, analogs, fragments, mimetics, mutants, synthetics, or variants thereof. This invention provides an isolated nucleic acid which encodes a human p-Hyde protein which induces susceptibility of a cancer cell to cell death, including allelic, analogs, fragments, mimetics, mutants, synthetics, or variants thereof.

15

20

25

30

35

In one embodiment the p-Hyde gene has a nucleotide sequence having at least 75% similarity with the nucleic acid coding sequence of SEQ ID NO: 1. In another embodiment the nucleic acid has a nucleotide sequence having at least 85% similarity with the nucleic acid coding sequence of SEQ ID NO: 1. In another embodiment the nucleic acid has a nucleotide sequence having at least 90% similarity with the nucleic acid coding sequence of SEQ ID NO: 1. In another embodiment the nucleic acid has a nucleotide sequence having at least 95% similarity with the nucleic acid coding sequence of SEQ ID NO: 1. In another embodiment the nucleic acid fragment comprises a fragment which begins at the nucleic acid at position 1 of the SEQ ID NO: 1 and ends at position 557 of SEQ ID NO: 1. In another embodiment the nucleic acid comprises a fragment which begins at the nucleic acid at position 1 of SEQ ID NO: 1 and ends at position 158 of SEQ ID NO: 1. In another embodiment the nucleic acid comprises a fragment which begins at the nucleic acid at position 50 of SEQ ID NO: 1 and ends at position 120 of SEQ ID NO: 1. The nucleic acid is DNA, cDNA, genomic DNA, or RNA.

TACGACTTGGTCAACCTGGCAGTCAAGCAGGTCTTGGCCAACAAGAGCCACC  
TCTGGGTGGAGGAGGAGGTCTGGCGGATGGAGATCTACCTCTCCCTGGGAGT  
GCTGGCCCTCGGCACGTTGTCCCTGCTGGCCGTGACCTCACTGCCGTCCATTG

5 CAAACTCGCTCAACTGGAGGGAGTTCAGCTTCGTTTCAGTCCTCACTGGGCTTT  
GTGGCCCTCGTGCTGAGCACACTGCACACGCTCACCTACGGCTGGACCCGCG  
CCTTCGAAGAGACCGCTACAAGTTCTACCTGCCTCCCACCTTCACGCTCACGC  
TGCTGGTGCCCTGCGTCGTCATCCTGGCCAAAGCCCTGTTTCTCCTGCCCTGC  
ATCAGCCGCAGACTCCCCAGGATCCGGAGAAGCTGGGAGAGGGAGAGCACC  
10 ATCAAGTTCACGCTGCCCACAGACCACGCCCTGGCCGAGAAGACGAGCCACG  
TATGAGGTGCCTGCCCTGGGCTCTGGACCCCGGGCACACGAGGGACGGTGCC  
CTGAGCCCGTTAGGTTTTCTTTTCTTGGTGGTGCAAAGTGGTATAACTGTGTG  
CAAATAGGAGGTTTGAGGTCCAAATTCCTGGGACTCAAATGTATGCATGACT  
ATTCAGAATGATATACACACATATGTGTATATGTATTTACATATATTCCACAT  
15 ATATAACAGGATTTGCAATTATACATAGCTAGCTAAAAAAAAAAAAAAAAAAAA  
(SEQ ID NO: 1)

In another embodiment the nucleic acid encodes an amino acid sequence having the sequence  
as set forth in SEQ ID NO 2. In one embodiment the the amino acid fragment comprises a  
20 fragment which begins at the amino acid at position 1 of SEQ ID NO 2 and ends at position  
101. In one embodiment the amino acid fragment comprises a fragment which begins at the  
amino acid at position 1 of SEQ ID NO 2 and ends at position 80. In one embodiment the  
amino acid fragment comprises a fragment which begins at the amino acid at position 1 of  
SEQ ID NO 2 and ends at position 60. In one embodiment the amino acid has at least 75%  
25 similarity with the nucleic acid coding sequence of SEQ ID NO 2. In another embodiment  
the amino acid has at least 85% similarity with the nucleic acid coding sequence of SEQ ID  
NO 2. In another embodiment the amino acid has at least 95% similarity with the nucleic  
acid coding sequence of SEQ ID NO 2.

30 MEIYLSLGVLALGTLSELLAVTSLPSIANSLNWREFSFVQSSLGFVALVLSTLHTLTY  
GWTRAFEETATSSSTCLPPSRSRCWCPASSSWPKPCFSCPASAADSPGSGEAGRGR  
APSSSRCPQTTPWPRRRATYEVPALGSGPRAHEGRCPEPVRFSFLGGAKWYNVCV  
QIGGLRSKFLGLKCMHDYSE (SEQ ID NO 2)

35 In one embodiment the p-Hyde gene has a nucleotide sequence having at least 75% similarity

5 with the nucleic acid coding sequence of SEQ ID NO: 3. In another embodiment the nucleic acid has a nucleotide sequence having at least 85% similarity with the nucleic acid coding sequence of SEQ ID NO: 3. In another embodiment the nucleic acid has a nucleotide sequence having at least 90% similarity with the nucleic acid coding sequence of SEQ ID NO: 3. In another embodiment the nucleic acid has a nucleotide sequence having at least 95% similarity with the nucleic acid coding sequence of SEQ ID NO: 3. In another embodiment the nucleic acid fragment comprises a fragment which begins at the nucleic acid at position 1 of the SEQ ID NO: 3 and ends at position 557 of SEQ ID NO: 3. In another embodiment the nucleic acid comprises a fragment which begins at the nucleic acid at position 1 of SEQ ID NO: 3 and ends at position 158 of SEQ ID NO: 3. In another embodiment the nucleic acid comprises a fragment which begins at the nucleic acid at position 50 of SEQ ID NO: 1 and ends at position 120 of SEQ ID NO: 3. The nucleic acid is DNA, cDNA, genomic DNA, or RNA.

20 ATGTCCGGGGAGATGGACAAACCGCTCATCAGTCGCCGCTTGGTGGACAGTG  
ATGGCAGTCTGGCTGAGGTCCCCAAGGAGGCTCCCAAAGTGGGCATCCTGGG  
CAGCGGGGATTTTGCCCGGTCCCTGGCCACACGCCTGGTGGGCTCTGGCTTCT  
TTGTGGTGGTGGGAAGCCGTAACCCCAAACGCACTGCCGGCCTCTTCCCCTCC  
TTAGCCCAAGTGACTTTCCAGGAGGAGGCCGTGAGCTCTCCAGAGGTCATCT  
TTGTGGCCGTGTTCCGGGAGCACTACTCCTCACTGTGCAGTCTTGCTGACCAG  
25 TTGGCTGGCAAGATCCTAGTGGATGTAAGCAACCCACGGAGAAAGGAGCGTC  
TTCAGCACCGCCAGTCGAACGCCGAGTACCTGGCCTCCCTCTTCCCTGCCTGC  
ACTGTGGTCAAGGCCTTCAACGTCATCTCTGCATGGGCCCTACAGGCTGGCCC  
AAGGGATGGGAACAGGCAGGTGCTCATCTGCGGTGACCAGCTGGAAGCCAA  
GCACACCGTCTCAGAGATGGCGCGCGCCATGGGTTTCACCCCACTGGACATG  
30 GGATCCCTGGCCTCAGCGAGGGAGGTAGAGGCCATAACCCCTGCGCCTCCTTC  
CATCCTGGAAGGTGCCCACCCTCCTGGCCCTGGGGCTAAGCACACAAAGCTA  
TGCCTACAACCTTCATCCGGGACGTTCTACAGCCGTACATCCGGAAAGATGAG  
AACAAGTTCTACAAGATGCCCCTGTCTGTGGTCAACACCACGaTACCCTGTGT  
GGCTTACGTGCTGCTGTCCCTGGTTTACCTGCCTGGTGTGCTGGCAGCTGCCC  
35 TTCAGCTGAGGAGGGGGACCAAGTACCAGCGCTTCCCAGACTGGCTGGACCA

5 TTGGCTGCAGCACCGCAAGCAGATCGGGCTACTCAGCTTTTTTTTCGCCATGC  
TGCACGCTCTCTACAGCTTCTGCCTGCCGCTGCGCCGCTCCCACCGCTATGAT  
CTGGTCAACCTGGCTGTGAAGCAGGTCCTGGCCAACAAGAGCCGCCTCTGGG  
TTGAGGAAGAAGTCTGGCGGATGGAGATATACCTGTCCCTGGGTGTGCTGGC  
TCTGGGCATGCTGTCACTGCTGGCGGTTACCTCGATCCCTTCCATTGCAAAC  
10 CACTCAACTGGAAGGAGTTCAGCTTTGTGCAGTCCACGCTGGGCTTCGTGGCC  
CTGATGCTGAGCACAATGCACACCCTCACCTACGGCTGGACCCGTGCTTTTGA  
GGAAAACCACTACAAGTTCTACCTGCCACCCACATTCACGCTCACGCTGCTCC  
TGCCCTGTGTCATCATCCTGGCCAAGGGCCTCTTCCTCCTGCCCTGCCTCAGC  
CACAGACTCACCAAGATCCGCAGGGGCTGGGAGAGGGATGGTGCCGTCAAG  
15 TTCATGCTGCCCCGCTGGCCACACACAGGGGGAGAAAACAAGCCACGTGTGA  
(SEQ ID NO: 3)

20 Further this invention provides an isolated nucleic acid which encodes an amino acid  
sequence NFIRDVLQPYIRKDENK. (SEQ ID NO: 4).

The nucleotide encoding p-Hyde includes RNA, cDNA, genomic DNA, synthetic forms, and  
mixed polymers, both sense and antisense strands, and may be chemically or  
biochemically modified or may contain non-natural or derivatized nucleotide bases, as will  
25 be readily appreciated by those skilled in the art. Such modifications include, for  
example, labels, methylation, substitution of one or more of the naturally occurring  
nucleotides with an analog, internucleotide modifications such as uncharged linkages (e.g.,  
methyl phosphonates, phosphotriesters, phosphoamidates, carbamates, etc.), charged  
linkages (e.g., phosphorothioates, phosphorodithioates, etc.), pendent moieties (e.g.,  
30 polypeptides), intercalators (e.g., acridine, psoralen, etc.), chelators, alkylators, and modified  
linkages (e.g., alpha anomeric nucleic acids, etc.). Also included are synthetic molecules that  
mimic nucleotides in their ability to bind to a designated sequence via hydrogen bonding  
and other chemical interactions. Such molecules are known in the art and include, for  
example, those in which peptide linkages substitute for phosphate linkages in the backbone  
35 of the molecule. substantially homologous to primary structural sequence but which include,

5 e.g., in vivo or in vitro chemical and biochemical modifications or which incorporate  
unusual amino acids. The nucleic acid may be modified. Such modifications include, for  
example, acetylation, carboxylation, phosphorylation, glycosylation, ubiquitination, labeling,  
e.g., with radionuclides, and various enzymatic modifications, as will be readily appreciated  
by those well skilled in the art. A variety of methods for labeling polypeptides and of  
10 substituents or labels useful for such purposes are well known in the art, and include  
radioactive isotopes such as  $^{32}\text{P}$ , ligands which bind to labeled antiligands (e.g.,  
antibodies), fluorophores, chemiluminescent agents, enzymes, and antiligands which  
can serve as specific binding pair members for a labeled ligand. The choice of label  
depends on the sensitivity required, ease of conjugation with the primer, stability  
15 requirements, and available instrumentation. Methods of labeling polypeptides are well  
known in the art. See, e.g., Sambrook et al., 1989 or Ausubel et al., 1992. Besides  
substantially full-length p-Hyde, the present invention provides for biologically active  
fragments of the p-hyde which are known to those skilled in the art.

20 As defined herein an "isolated" or "substantially pure" nucleic acid (e.g., an RNA, DNA or  
a mixed polymer) is one which is substantially separated from other cellular  
components which naturally accompany a native human sequence or protein, e.g.,  
ribosomes, polymerases, many other human genome sequences and proteins. The term  
embraces a nucleic acid sequence or protein which has been removed from its naturally  
25 occurring environment, and includes recombinant or cloned DNA isolates and chemically  
synthesized analogs or analogs biologically synthesized by heterologous systems.

"p-Hyde Allele" refers to normal alleles of the p-Hyde locus as well as alleles carrying  
variations that predispose individuals to develop cancer of many sites including, for  
30 example, breast, ovarian, colorectal and prostate cancer. Such predisposing alleles are  
also called "p-Hyde susceptibility alleles".

"p-Hyde Locus," "p-Hyde Gene," "p-Hyde Nucleic Acids" or "p-Hyde  
Polynucleotide" each refer to polynucleotides, all of which are in the p-Hyde region, that  
35 are likely to be expressed in normal tissue, certain alleles of which predispose an



5 individual to develop breast, ovarian, colorectal and prostate cancers. Mutations at the  
p-Hyde locus may be involved in the initiation and/or progression of other types of tumors.  
The locus is indicated in part by mutations that predispose individuals to develop cancer.  
These mutations fall within the p-Hyde region described infra. The p-Hyde locus is  
intended to include coding sequences, intervening sequences and regulatory elements  
10 controlling transcription and/or translation. The p-Hyde locus is intended to include all  
allelic variations of the DNA sequence.

A "nucleic acid" refers to the phosphate ester polymeric form of ribonucleosides (adenosine,  
guanosine, uridine or cytidine; "RNA molecules") or deoxyribonucleosides (deoxyadenosine,  
15 deoxyguanosine, deoxythymidine, or deoxycytidine; "DNA molecules") in either single  
stranded form, or a double-stranded helix. Double stranded DNA-DNA, DNA-RNA and  
RNA-RNA helices are possible. The term nucleic acid molecule, and in particular DNA or  
RNA molecule, refers only to the primary and secondary structure of the molecule, and does  
not limit it to any particular tertiary forms. Thus, this term includes double-stranded DNA  
20 found, *inter alia*, in linear or circular DNA molecules (*e.g.*, restriction fragments), plasmids,  
and chromosomes. In discussing the structure of particular double-stranded DNA molecules,  
sequences may be described herein according to the normal convention of giving only the  
sequence in the 5' to 3' direction along the nontranscribed strand of DNA (*i.e.*, the strand  
having a sequence homologous to the mRNA). A "recombinant DNA" is a DNA that has  
25 undergone a molecular biological manipulation.

The phrase "nucleic acid encoding" refers to a nucleic acid molecule which directs the  
expression of a specific protein or peptide. The nucleic acid sequences include both the  
DNA strand sequence that is transcribed into RNA and the RNA sequence that is translated  
30 into protein. The nucleic acid molecule include both the full length nucleic acid sequences  
as well as non-full length sequences derived from the full length protein. It being further  
understood that the sequence includes the degenerate codons of the native sequence or  
sequences which may be introduced to provide codon preference in a specific host cell

35 "Recombinant nucleic acid" is a nucleic acid which is not naturally occurring, or which

5 is made by the artificial combination of two otherwise separated segments of sequence.  
This artificial combination is often accomplished by either chemical synthesis means,  
or by the artificial manipulation of isolated segments of nucleic acids, e.g., by genetic  
engineering techniques. Such is usually done to replace a codon with a redundant codon  
encoding the same or a conservative amino acid, while typically introducing or  
10 removing a sequence recognition site. Alternatively, it is performed to join together  
nucleic acid segments of desired functions to generate a desired combination of functions.

As used herein, the term "cancer cell" means a tissue that grows by cellular proliferation  
more rapidly than normal, e.g., more rapidly than adjoining cells, or other cells in the tissue.  
15 Neoplastic cells continue to grow after growth stimuli cease. Generally, tumors represent  
or form a distinct mass of tissue. The present invention relates to both types of tumors, but  
is particularly valuable in the treatment of cancers.

In one embodiment the cancer cells are selected from a group consisting of: melanoma;  
20 lymphoma; leukemia; and prostate, colorectal, pancreatic, breast, brain, or gastric  
carcinoma. Examples of tumors include but are not limited to: include sarcomas and  
carcinomas such as, but not limited to: fibrosarcoma, myxosarcoma, liposarcoma,  
chondrosarcoma, osteogenic sarcoma, chordoma, angiosarcoma, endotheliosarcoma,  
lymphangiosarcoma, lymphangioendotheliosarcoma, synovioma, mesothelioma, Ewing's  
25 sarcoma, leiomyosarcoma, rhabdomyosarcoma, colon carcinoma, pancreatic cancer, breast  
cancer, ovarian cancer, prostate cancer, squamous cell carcinoma, basal cell carcinoma,  
adenocarcinoma, sweat gland carcinoma, sebaceous gland carcinoma, papillary carcinoma,  
papillary adenocarcinomas, cystadenocarcinoma, medullary carcinoma, bronchogenic  
carcinoma, renal cell carcinoma, hepatoma, bile duct carcinoma, choriocarcinoma,  
30 seminoma, embryonal carcinoma, Wilms' tumor, cervical cancer, germ tumor, non-small cell  
lung carcinoma, small cell lung carcinoma, bladder carcinoma, epithelial carcinoma, glioma,  
astrocytoma, medulloblastoma, craniopharyngioma, ependymoma, pinealoma,  
hemangioblastoma, acoustic neuroma, oligodendroglioma, meningioma, melanoma,  
neuroblastoma, and retinoblastoma. In a preferred embodiment the tumor is a melanoma or  
35 a prostate cell.

5 Mutations can be made in a nucleic acid encoding p-Hyde such that a particular codon is  
changed to a codon which codes for a different amino acid but the induction of susceptibility  
to cell death is maintained. Such a mutation is generally made by making the fewest  
nucleotide changes possible. A substitution mutation of this sort can be made to change an  
10 amino acid in the resulting protein in a non-conservative manner (i.e., by changing the codon  
from an amino acid belonging to a grouping of amino acids having a particular size or  
characteristic to an amino acid belonging to another grouping) or in a conservative manner  
(i.e., by changing the codon from an amino acid belonging to a grouping of amino acids  
having a particular size or characteristic to an amino acid belonging to the same grouping).  
Such a conservative change generally leads to less change in the structure and function of the  
15 resulting protein. A non-conservative change is more likely to alter the structure, activity or  
function of the resulting protein. The present invention should be considered to include  
sequences containing conservative changes which do not significantly alter the activity or  
binding characteristics of the resulting protein. Substitutes for an amino acid within the  
sequence may be selected from other members of the class to which the amino acid belongs.  
20 For example, the nonpolar (hydrophobic) amino acids include alanine, leucine, isoleucine,  
valine, proline, phenylalanine, tryptophan and methionine. Amino acids containing aromatic  
ring structures are phenylalanine, tryptophan, and tyrosine. The polar neutral amino acids  
include glycine, serine, threonine, cysteine, tyrosine, asparagine, and glutamine. The  
positively charged (basic) amino acids include arginine, lysine and histidine. The negatively  
25 charged (acidic) amino acids include aspartic acid and glutamic acid. Such alterations will  
not be expected to affect apparent molecular weight as determined by polyacrylamide gel  
electrophoresis, or isoelectric point. This isolated nucleic acid also encodes mutant p-Hyde  
or the wildtype protein.

30 This invention provides for a replicable vector comprising the isolated nucleic acid molecule  
of the DNA virus. The vector includes, but is not limited to: a plasmid, cosmid, phage or  
yeast artificial chromosome (YAC) which contains at least a portion of the isolated nucleic  
acid molecule. As an example to obtain these vectors, insert and vector DNA can both be  
exposed to a restriction enzyme to create complementary ends on both molecules which base  
35 pair with each other and are then ligated together with DNA ligase. Alternatively, linkers

5 can be ligated to the insert DNA which correspond to a restriction site in the vector DNA,  
which is then digested with the restriction enzyme which cuts at that site. Other means are  
also available and known to an ordinary skilled practitioner. In one embodiment the  
adenovirus vector is a replication-deficient adenovirus type 5 expression vector. In another  
10 embodiment the adenovirus vector comprises an adenovirus genome having a deletion in the  
E1 and E3 region of the genome and an insertion within the region of a nucleic acid encoding  
p-Hyde under the control of a promoter. The promoter may be a Rous Sarcoma virus  
promoter.

15 Knowledge of the genetic organization of adenovirus, a 36 kB, linear and double-stranded  
DNA virus, allows substitution of a large piece of adenoviral DNA with foreign sequences  
up to 7 kB (Grunhaus and Horwitz, 1992). In contrast to retrovirus, the infection of  
adenoviral DNA into host cells does not result in chromosomal integration because  
adenoviral DNA can replicate in an episomal manner without potential genotoxicity. Also,  
adenoviruses are structurally stable, and no genome rearrangement has been detected after  
20 extensive amplification. Adenovirus can infect virtually all epithelial cells regardless of their  
cell cycle stage. So far, adenoviral infection appears to be linked only to mild disease such  
as acute respiratory disease in the human.

25 Adenovirus is particularly suitable for use as a gene transfer vector because of its mid-sized  
genome, ease of manipulation, high titer, wide target-cell range, and high infectivity. Both  
ends of the viral genome contain 100-200 base pair Cop) inverted terminal repeats (ITR),  
which are cis elements necessary for viral DNA replication and packaging. The early (E) and  
late (L) regions of the genome contain different transcription units that are divided by the  
onset of viral DNA replication. The E1 region (E1A and E1B) encodes proteins responsible  
30 for the regulation of transcription of the viral genome and a few cellular genes. The  
expression of the E2 region (E2A and E2B) results in the synthesis of the proteins for viral  
DNA replication. These proteins are involved in DNA replication, late gene expression, and  
host cell shut off (Renan, 1990). The products of the late genes, including the majority of the  
viral capsid proteins, are expressed only after significant processing of a single primary  
35 transcript issued by the major late promoter (MLP). The MLP (located at 16.8 m.u.) is

5 particularly efficient during the late phase of infection, and all the mRNAs issued from this promoter possess a 5' tripartite leader (TL) sequence which makes them preferred mRNAs for translation.

10 In the current system, recombinant adenovirus is generated from homologous recombination between shuttle vector and provirus vector. Due to the possible recombination between two proviral vectors, wild-type adenovirus may be generated from this process. Therefore, it is critical to isolate a single clone of virus from an individual plaque and examine its genomic structure. Use of the YAC system is an alternative approach for the production of recombinant adenovirus.

15 Generation and propagation of the current adenovirus vectors, which are replication deficient, depend on a unique helper cell line, designated 293, which was transformed from human embryonic kidney cells by Ad5 DNA fragments and constitutively expresses E1 proteins (Graham, et al., 1977). Since the E3 region is dispensable from the adenovirus genome (Jones and Shenk, 1978), the current adenovirus vectors, with the help of 293 cells, carry foreign DNA in either the E1, the E3 or both regions (Graham and Prevec, 1991). In nature, adenovirus can package approximately 105% of the wild-type genome (Ghosh-Choudhury, et al., 1987), providing capacity for about 2 extra kB of DNA. Combined with the approximately 5.5 kB of DNA that is replaceable in the E1 and E3 regions, the maximum capacity of the current adenovirus vector is under 7.5 kB, or about 15% of the total length of the vector. More than 80% of the adenovirus viral genome remains in the vector backbone and is the source of vector-borne cytotoxicity. Also, the replication deficiency of the E1 deleted virus is incomplete. For example, leakage of viral gene expression has been observed with the currently available adenovirus vectors at high multiplicities of infection (Mulligan, 25 30 1993).

Helper cell lines may be derived from human cells such as human embryonic kidney cells, muscle cells, hematopoietic cells or other human embryonic mesenchymal or epithelial cells. Alternatively, the helper cells may be derived from the cells of other mammalian species that are permissive for human adenovirus. Such cells include, e.g., Vero cells or other monkey 35



5 embryonic mesenchymal or epithelial cells. As stated above, the preferred helper cell line is 293.

10 Other than the requirement that the adenovirus vector be replication defective, or at least conditionally defective, the nature of the adenovirus vector is not believed to be crucial to the successful practice of the invention. The adenovirus may be of any of the 42 different known serotypes or subgroups A-F. Adenovirus type 5 of subgroup C is the preferred starting material in order to obtain the conditional replication-defective adenovirus vector for use in the method of the present invention. This is because Adenovirus type 5 is a human adenovirus about which a great deal of biochemical and genetic information is known, and 15 it has historically been used for most constructions employing adenovirus as a vector.

20 As stated above, the typical vector according to the present invention is replication defective and will not have an adenovirus E1 region. Thus, it will be most convenient to introduce the nucleic acid encoding p-Hyde at the position from which the E1 coding sequences have been removed. However, the position of insertion of the p-Hyde coding region within the adenovirus sequences is not critical to the present invention. The nucleic acid encoding a p-Hyde transcription unit also may be inserted in lieu of the deleted E3 region in E3 replacement vectors as described previously by Karlsson et. al. (1986) or in the E4 region where a helper cell line or helper virus complements the E4 defect. 25

30 Adenovirus is easy to grow and manipulate and exhibits broad host range in vitro and in vivo. This group of viruses can be obtained in high titers, e.g.,  $10^{9-11}$  plaque-forming unit (PFU)/ml, and they are highly infective. The life cycle of adenovirus does not require integration into the host cell genome. The foreign genes delivered by adenovirus vectors are episomal, and therefore, have low genotoxicity to host cells. No side effects have been reported in studies of vaccination with wild-type adenovirus (Couch et al., 1963; Top et al., 1971), demonstrating their safety and therapeutic potential as in vivo gene transfer vectors.

35 Adenovirus vectors have been used in eukaryotic gene expression (Levrero et al., 1991;

5 Gomez-Foix et al., 1992) and vaccine development (Grunhaus and Horwitz, 1992; Graham  
and Prevec, 1992). Recently, animal studies suggested that recombinant adenovirus could  
be used for gene therapy (Stratford-Perricaudet and Perricaudet, 1991; Stratford-Perricaudet  
et al., 1990; Rich et al., 1993). Experiments in administering recombinant adenovirus to  
10 different tissues include trachea instillation (Rosenfeld et al., 1991; Rosenfeld et al., 1992),  
muscle injection (Ragot et al., 1993), peripheral intravenous injection (Herz and Gerard,  
1993), and stereotactic inoculation into the brain (Le Gal La Salle et al., 1993).

15 An appropriate promoter and other necessary vector sequences will be selected so as to  
be functional in the host, and may include, when appropriate, those naturally associated  
with p-Hyde genes. Examples of workable combinations of cell lines and expression vectors  
are described in Sambrook et al., 1989 or Ausubel et al., 1992; see also, e.g., Metzger et al.,  
1988. Many useful vectors are known in the art and may be obtained from such vendors  
as Stratagene, New England Biolabs, Promega Biotech, and others. Promoters such as the  
trp, lac and phage promoters, tRNA promoters and glycolytic enzyme promoters may be  
20 used in prokaryotic hosts. Useful yeast promoters include promoter regions for  
metallothionein, 3-phosphoglycerate kinase or other glycolytic enzymes such as enolase  
or glyceraldehyde-3-phosphate dehydrogenase, enzymes responsible for maltose and  
galactose utilization, and others. Vectors and promoters suitable for use in yeast  
expression are further described in Hitzeman et al., EP 73,675A. Appropriate non-native  
25 mammalian promoters might include the early and late promoters from SV40 (Fiers et al.,  
1978) or promoters derived from murine Moloney leukemia virus, mouse minor virus,  
avian sarcoma viruses, adenovirus II, bovine papilloma virus or polyoma. In addition,  
the construct may be joined to an amplifiable gene (e.g., DHFR) so that multiple copies  
of the gene may be made. For appropriate enhancer and other expression control sequences,  
30 see also Enhancers and Eukaryotic Gene Expression, Cold Spring Harbor Press, Cold  
Spring Harbor, N.Y. (1983).

While such expression vectors may replicate autonomously, they may also replicate by  
being inserted into the genome of the host cell, by methods well known in the art.  
35 Expression and cloning vectors will likely contain a selectable marker, a gene encoding a

5 protein necessary for survival or growth of a host cell transformed with the vector. The presence of this gene ensures growth of only those host cells which express the inserts. Typical selection genes encode proteins that a) confer resistance to antibiotics or other toxic substances, e.g. ampicillin, neomycin, methotrexate, etc.; b) complement auxotrophic deficiencies, or c) supply critical nutrients not available from complex media, e.g., the gene  
10 encoding D-alanine racemase for Bacilli. The choice of the proper selectable marker will depend on the host cell, and appropriate markers for different hosts are well known in the art.

The vectors containing the nucleic acids of interest can be transcribed in vitro, and the resulting RNA introduced into the host cell by well-known methods, e.g., by injection (see,  
15 Kubo et al., 1988), or the vectors can be introduced directly into host cells by methods well known in the art, which vary depending on the type of cellular host, including electroporation; transfection employing calcium chloride, rubidium chloride, calcium phosphate, DEAE-dextran, or other substances; microprojectile bombardment; lipofection; infection (where the vector is an infectious agent, such as a retroviral genome); and other  
20 methods. See generally, Sambrook et al., 1989 and Ausubel et al., 1992. The introduction of the polynucleotides into the host cell by any method known in the art, including, inter alia, those described above, will be referred to herein as "transformation." The cells into which have been introduced nucleic acids described above are meant to also include the progeny of such cells.

25 Regulatory elements required for expression include promoter or enhancer sequences to bind RNA polymerase and transcription initiation sequences for ribosome binding. For example, a bacterial expression vector includes a promoter such as the lac promoter and for transcription initiation the Shine-Dalgarno sequence and the start codon AUG. Similarly,  
30 a eukaryotic expression vector includes a heterologous or homologous promoter for RNA polymerase II, a downstream polyadenylation signal, the start codon AUG, and a termination codon for detachment of the ribosome. Such vectors may be obtained commercially or assembled from the sequences described by methods well-known in the art, for example the methods described above for constructing vectors in general.

35

5       Viral promoters, cellular promoters/enhancers and inducible promoters/enhancers that could  
be used in combination with the nucleic acid encoding a p-Hyde in an expression construct  
include but are not limited to the following: Immunoglobulin Heavy Chain;  
Immunoglobulin Light Chain; T-Cell Receptor; HLA DQ alpha and DQ beta; beta -  
Interferon; Interleukin-2; Interleukin-2 Receptor; MHC Class II 5 alpha; MHC Class II HLA-  
10       DR alpha; beta -Actin; Muscle Creatine Kinase; Prealbumin (Transthyretin); Elastase I;  
Metallothionein; Collagenase; Albumin Gene; alpha -Fetoprotein; tau -Globin; beta -Globin;  
c-fos; c-HA-ras; Neural Cell Adhesion Molecule (NCAM); alpha 1-Antitrypsin; H2B  
(TH2B) Histone; Mouse or Type I Collagen; Glucose-Regulated Proteins (GRP94 and  
GRP78); Rat Growth Hormon; Human Serum Amyloid A (SAA); Troponin I (TN I);  
15       Platelet-Derived Growth Factor; Duchenne Muscular; SV40; Polyoma; Retroviruses;  
Papilloma Virus; Hepatitis B Virus; Human Immunodeficiency Virus; Cytomegalovirus;  
Gibbon Ape Leukemia Virus; MT II; MMTV (mouse mammary       Glucocorticoids;  
Adenovirus 5 E2.

20       Additionally any promoter/enhancer combination (as per the Eukaryotic Promoter Data Base  
EPDB) could also be used to drive expression of a p-Hyde. Use of a T3, T7 or SP6  
cytoplasmic expression system is another possible embodiment. Eukaryotic cells can support  
cytoplasmic transcription from certain bacterial promoters if the appropriate bacterial  
polymerase is provided, either as part of the delivery complex or as an additional genetic  
25       expression construct.

This invention provides a host cell containing the above vector. The host cell may contain  
the isolated DNA molecule artificially introduced into the host cell. The host cell may be a  
eukaryotic or bacterial cell (such as E.coli), yeast cells, fungal cells, insect cells and animal  
30       cells. Suitable animal cells include, but are not limited to Vero cells, HeLa cells, Cos cells,  
CV1 cells and various primary mammalian cells.

The term "vector", refers to viral expression systems, autonomous self-replicating circular  
DNA (plasmids), and includes both expression and nonexpression plasmids. Where a  
35       recombinant microorganism or cell culture is described as hosting an "expression vector,"

5 this includes both extrachromosomal circular DNA and DNA that has been incorporated into the host chromosome(s). Where a vector is being maintained by a host cell, the vector may either be stably replicated by the cells during mitosis as an autonomous structure, or is incorporated within the host's genome.

10 The term "plasmid" refers to an autonomous circular DNA molecule capable of replication in a cell, and includes both the expression and nonexpression types. Where a recombinant microorganism or cell culture is described as hosting an "expression plasmid", this includes latent viral DNA integrated into the host chromosome(s). Where a plasmid is being maintained by a host cell, the plasmid is either being stably replicated by the cells during  
15 mitosis as an autonomous structure or is incorporated within the host's genome.

The following terms are used to describe the sequence relationships between two or more nucleic acid molecules or polynucleotides: "reference sequence", "comparison window", "sequence identity", "percentage of sequence identity", and "substantial identity". A  
20 "reference sequence" is a defined sequence used as a basis for a sequence comparison; a reference sequence may be a subset of a larger sequence, for example, as a segment of a full-length cDNA or gene sequence given in a sequence listing or may comprise a complete cDNA or gene sequence.

25 Optimal alignment of sequences for aligning a comparison window may be conducted by the local homology algorithm of Smith and Waterman (1981) *Adv. Appl. Math.* 2:482, by the homology alignment algorithm of Needleman and Wunsch (1970) *J. Mol. Biol.* 48:443, by the search for similarity method of Pearson and Lipman (1988) *Proc. Natl. Acad. Sci. (USA)* 85:2444, or by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package Release 7.0, Genetics Computer  
30 Group, 575 Science Dr., Madison, WI).

"Substantial identity" or "substantial sequence identity" mean that two peptide sequences, when optimally aligned, such as by the programs GAP or BESTFIT using default gap which  
35 share at least 90 percent sequence identity, preferably at least 95 percent sequence identity,



5 more preferably at least 99 percent sequence identity or more. "Percentage amino acid identity" or "percentage amino acid sequence identity" refers to a comparison of the amino acids of two polypeptides which, when optimally aligned, have approximately the designated percentage of the same amino acids. For example, "95% amino acid identity" refers to a comparison of the amino acids of two polypeptides which when optimally aligned have 95% amino acid identity. Preferably, residue positions which are not identical differ by conservative amino acid substitutions. For example, the substitution of amino acids having similar chemical properties such as charge or polarity are not likely to effect the properties of a protein. Examples include glutamine for asparagine or glutamic acid for aspartic acid.

15 This invention provides a nucleic acid having a sequence complementary to the sequence of the isolated nucleic acid of the human p-Hyde gene. Specifically, this invention provides an oligonucleotide of at least 15 nucleotides capable of specifically hybridizing with a sequence of nucleotides present within a nucleic acid which encodes the human p-Hyde. In one embodiment the nucleic acid is DNA or RNA. In another embodiment the oligonucleotide is labeled with a detectable marker. In another embodiment the oligonucleotide is a radioactive isotope, a fluorophor or an enzyme.

Oligonucleotides which are complementary may be obtained as follows: The polymerase chain reaction is then carried out using the two primers. See *PCR Protocols: A Guide to Methods and Applications* [74]. Following PCR amplification, the PCR-amplified regions of a viral DNA can be tested for their ability to hybridize to the three specific nucleic acid probes listed above. Alternatively, hybridization of a viral DNA to the above nucleic acid probes can be performed by a Southern blot procedure without viral DNA amplification and under stringent hybridization conditions as described herein.

30 Oligonucleotides for use as probes or PCR primers are chemically synthesized according to the solid phase phosphoramidite triester method first described by Beaucage and Carruthers [19] using an automated synthesizer, as described in Needham-VanDevanter [69]. Purification of oligonucleotides is by either native acrylamide gel electrophoresis or by anion-exchange HPLC as described in Pearson, J.D. and Regnier, F.E. [75A]. The sequence

5 of the synthetic oligonucleotide can be verified using the chemical degradation method of Maxam, A.M. and Gilbert, W. [63].

10 High stringent hybridization conditions are selected at about 5° C lower than the thermal melting point ( $T_m$ ) for the specific sequence at a defined ionic strength and pH. The  $T_m$  is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. Typically, stringent conditions will be those in which the salt concentration is at least about 0.02 molar at pH 7 and the temperature is at least about 60°C. As other factors may significantly affect the stringency of hybridization, including, among others, base composition and size of the complementary strands, the presence of organic solvents, ie. salt or formamide concentration, and the extent of base mismatching, the combination of parameters is more important than the absolute measure of any one. For Example high stringency may be attained for example by overnight hybridization at about 68°C in a 6x SSC solution, washing at room temperature with 6x SSC solution, followed by washing at about 68°C in a 6x SSC in a 0.6x SSX solution.

20 Hybridization with moderate stringency may be attained for example by: 1) filter pre-hybridizing and hybridizing with a solution of 3x sodium chloride, sodium citrate (SSC), 50% formamide, 0.1M Tris buffer at Ph 7.5, 5x Denhardt's solution; 2.) pre-hybridization at 37°C for 4 hours; 3) hybridization at 37°C with amount of labelled probe equal to 3,000,000 cpm total for 16 hours; 4) wash in 2x SSC and 0.1% SDS solution; 5) wash 4x for 1 minute each at room temperature at 4x at 60°C for 30 minutes each; and 6) dry and expose to film.

30 The phrase "selectively hybridizing to" refers to a nucleic acid probe that hybridizes, duplexes or binds only to a particular target DNA or RNA sequence when the target sequences are present in a preparation of total cellular DNA or RNA. By selectively hybridizing it is meant that a probe binds to a given target in a manner that is detectable in a different manner from non-target sequence under high stringency conditions of hybridization. in a different "Complementary" or "target" nucleic acid sequences refer to those nucleic acid sequences which selectively hybridize to a nucleic acid probe. Proper

5 annealing conditions depend, for example, upon a probe's length, base composition, and the number of mismatches and their position on the probe, and must often be determined empirically. For discussions of nucleic acid probe design and annealing conditions, see, for example, Sambrook *et al.*, [81] or Ausubel, F., *et al.*, [8].

10 It will be readily understood by those skilled in the art and it is intended here, that when reference is made to particular sequence listings, such reference includes sequences which substantially correspond to its complementary sequence and those described including allowances for minor sequencing errors, single base changes, deletions, substitutions and the like, such that any such sequence variation corresponds to the nucleic acid sequence of the  
15 pathogenic organism or disease marker to which the relevant sequence listing relates.

The primer pairs of the present invention are useful for determination of the nucleotide sequence of a particular p-Hyde allele using PCR. The pairs of single-stranded DNA primers can be annealed to sequences within or surrounding the p-Hyde gene in order to  
20 prime amplifying DNA synthesis of the p-Hyde gene itself. A complete set of these primers allows synthesis of all of the nucleotides of the p-Hyde gene coding sequences, i.e., the exons. The set of primers preferably allows synthesis of both intron and exon sequences. Allele-specific primers can also be used. Such primers anneal only to particular p-Hyde mutant alleles, and thus will only amplify a product in the presence of the mutant allele  
25 as a template.

Nucleic acid probe technology is well known to those skilled in the art who readily appreciate that such probes may vary greatly in length and may be labeled with a detectable label, such as a radioisotope or fluorescent dye, to facilitate detection of the probe. DNA  
30 probe molecules may be produced by insertion of a DNA molecule having the full-length or a fragment of the isolated nucleic acid molecule of the DNA virus into suitable vectors, such as plasmids or bacteriophages, followed by transforming into suitable bacterial host cells, replication in the transformed bacterial host cells and harvesting of the DNA probes, using methods well known in the art. Alternatively, probes may be generated chemically from  
35 DNA synthesizers.

5 RNA probes may be generated by inserting the full length or a fragment of the isolated  
nucleic acid molecule of the DNA virus downstream of a bacteriophage promoter such as T3,  
T7 or SP6. Large amounts of RNA probe may be produced by incubating the labeled  
nucleotides with a linearized isolated nucleic acid molecule of the DNA virus or its fragment  
10 where it contains an upstream promoter in the presence of the appropriate RNA polymerase.

As defined herein nucleic acid probes may be DNA or RNA fragments. DNA fragments can  
be prepared, for example, by digesting plasmid DNA, or by use of PCR, or synthesized by  
either the phosphoramidite method described by Beaucage and Carruthers, [19], or by the  
triesther method according to Matteucci, *et al.*, [62], both incorporated herein by reference.  
15 A double stranded fragment may then be obtained, if desired, by annealing the chemically  
synthesized single strands together under appropriate conditions or by synthesizing the  
complementary strand using DNA polymerase with an appropriate primer sequence. Where  
a specific sequence for a nucleic acid probe is given, it is understood that the complementary  
strand is also identified and included. The complementary strand will work equally well in  
20 situations where the target is a double-stranded nucleic acid. It is also understood that when  
a specific sequence is identified for use a nucleic probe, a subsequence of the listed sequence  
which is 25 basepairs or more in length is also encompassed for use as a probe.

The nucleic acid of the subject invention also include DNA molecules coding for polypeptide  
25 analogs, fragments or derivatives of antigenic polypeptides which differ from naturally-  
occurring forms in terms of the identity or location of one or more amino acid residues  
(deletion analogs containing less than all of the residues specified for the protein, substitution  
analog wherein one or more residues specified are replaced by other residues and addition  
analog where in one or more amino acid residues is added to a terminal or medial portion  
30 of the polypeptides) and which share some or all properties of naturally-occurring forms.  
These molecules include: the incorporation of codons "preferred" for expression by selected  
non-mammalian hosts; the provision of sites for cleavage by restriction endonuclease  
enzymes; and the provision of additional initial, terminal or intermediate DNA sequences  
35 that facilitate construction of readily expressed vectors.

5 Also, this invention provides an antisense molecule capable of specifically hybridizing with  
the isolated nucleic acid of the human p-Hyde gene. This invention provides an antagonist  
capable of blocking the expression of the peptide or polypeptide encoded by the isolated  
DNA molecule. In one embodiment the antagonist is capable of hybridizing with a double  
10 stranded DNA molecule. In another embodiment the antagonist is a triplex oligonucleotide  
capable of hybridizing to the DNA molecule. In another embodiment the triplex  
oligonucleotide is capable of binding to at least a portion of the isolated DNA molecule with  
a nucleotide sequence..

15 The antisense molecule may be DNA or RNA or variants thereof (i.e. DNA or RNA with a  
protein backbone). The present invention extends to the preparation of antisense nucleotides  
and ribozymes that may be used to interfere with the expression of the receptor recognition  
proteins at the translation of a specific mRNA, either by masking that MRNA with an  
antisense nucleic acid or cleaving it with a ribozyme.

20 Antisense nucleic acids are DNA or RNA molecules that are complementary to at least a  
portion of a specific MRNA molecule. In the cell, they hybridize to that MRNA, forming  
a double stranded molecule. The cell does not translate an MRNA in this double-stranded  
form. Therefore, antisense nucleic acids interfere with the expression of MRNA into protein.

25 Antisense nucleotides or polynucleotide sequences are useful in preventing or  
diminishing the expression of the p-Hyde gene, as will be appreciated by those skilled in  
the art. For example, polynucleotide vectors containing all or a portion of the p-Hyde gene  
or other sequences from the p-Hyde region (particularly those flanking the p-Hyde gene)  
may be placed under the control of a promoter in an antisense orientation and introduced into  
30 a cell. Expression of such an antisense construct within a cell will interfere with p-Hyde  
transcription and/or translation and/or replication. Oligomers of about fifteen nucleotides and  
molecules that hybridize to the AUG initiation codon are particularly efficient, since they are  
easy to synthesize and are likely to pose fewer problems than larger molecules upon  
introduction to cells.

35 This invention provides a transgenic nonhuman mammal which comprises at least a portion



5 of the isolated DNA molecule introduced into the mammal at an embryonic stage. Methods of producing a transgenic nonhuman mammal are known to those skilled in the art.

10 This invention provides a polypeptide comprising the amino acid sequence of a human p-Hyde. In one embodiment the amino acid sequence is set forth in SEQ ID NO. 2. This invention provides a fusion protein or chimeric comprising the polypeptide. This invention provides an antibody which specifically binds to the polypeptide. In one embodiment the antibody is a monoclonal or polyclonal antibody.

15 The present invention further provides methods of preparing a polynucleotide comprising polymerizing nucleotides to yield a sequence comprised of at least eight consecutive nucleotides of the p-Hyde locus; and methods of preparing a polypeptide comprising polymerizing amino acids to yield a sequence comprising at least five amino acids encoded within the p-Hyde locus.

20 The present invention provides an isolated polynucleotide comprising all, or a portion of the p-Hyde locus or of a mutated p-Hyde locus. Such polynucleotides may be antisense polynucleotides. The present invention also provides a recombinant construct comprising such an isolated polynucleotide, for example, a recombinant construct suitable for expression in a transformed host cell.

25 Also provided by the present invention are methods of detecting a polynucleotide comprising a portion of the p-Hyde locus or its expression product in an analyte. Such methods may further comprise the step of amplifying the portion of the p-Hyde locus, and may further include a step of providing a set of polynucleotides which are primers for  
30 amplification of said portion of the p-Hyde locus. The method is useful for either diagnosis of the predisposition to cancer or the diagnosis or prognosis of cancer.

This invention also provides a method of producing a polypeptide encoded by isolated DNA molecule, which comprises growing the above host vector system under suitable conditions  
35 permitting production of the polypeptide and recovering the polypeptide so produced.

5 Further, the isolated polypeptide encoded by the isolated DNA molecule may be linked to a second polypeptide encoded by a nucleic acid molecule to form a fusion protein by expression in a suitable host cell. In one embodiment the second nucleic acid molecule encodes beta-galactosidase. Other nucleic acid molecules which are used to form a fusion protein are known to those skilled in the art.

10

This invention provides an antibody which specifically binds to the polypeptide encoded by the isolated DNA molecule. In one embodiment the antibody is a monoclonal antibody. In another embodiment the antibody is a polyclonal antibody. The antibody or DNA molecule may be labelled with a detectable marker including, but not limited to: a radioactive label, 15 or a colorimetric, a luminescent, or a fluorescent marker, or gold. Radioactive labels include, but are not limited to:  $^3\text{H}$ ,  $^{14}\text{C}$ ,  $^{32}\text{P}$ ,  $^{33}\text{P}$ ,  $^{35}\text{S}$ ,  $^{36}\text{Cl}$ ,  $^{51}\text{Cr}$ ,  $^{57}\text{Co}$ ,  $^{59}\text{Co}$ ,  $^{59}\text{Fe}$ ,  $^{90}\text{Y}$ ,  $^{125}\text{I}$ ,  $^{131}\text{I}$ , and  $^{186}\text{Re}$ . Fluorescent markers include but are not limited to: fluorescein, rhodamine and auramine. Colorimetric markers include, but are not limited to: biotin, and digoxigenin. Methods of producing the polyclonal or monoclonal antibody are known to those of ordinary skill in the 20 art.

20

Further, the antibody or nucleic acid molecule complex may be detected by a second antibody which may be linked to an enzyme, such as alkaline phosphatase or horseradish peroxidase. Other enzymes which may be employed are well known to one of ordinary skill 25 in the art.

25

"Specifically binds to an antibody" or "specifically immunoreactive with", when referring to a protein or peptide, refers to a binding reaction which is determinative of the presence of the p-Hyde of the invention in the presence of a heterogeneous population of proteins and 30 other biologics including viruses other than the p-Hyde. Thus, under designated immunoassay conditions, the specified antibodies bind to the p-Hyde antigens and do not bind in a significant amount to other antigens present in the sample. Specific binding to an antibody under such conditions may require an antibody that is selected for its specificity for a particular protein. For example, antibodies raised to the human p-Hyde immunogen 35 described herein can be selected to obtain antibodies specifically immunoreactive with the

35

5 p-Hyde proteins and not with other proteins. These antibodies recognize proteins homologous to the human p-Hyde protein. A variety of immunoassay formats may be used to select antibodies specifically immunoreactive with a particular protein. For example, solid-phase ELISA immunoassays are routinely used to select monoclonal antibodies specifically immunoreactive with a protein. See Harlow and Lane [32] for a description of  
10 immunoassay formats and conditions that can be used to determine specific immunoreactivity.

This invention provides a method to select specific regions on the polypeptide encoded by the isolated DNA molecule of the DNA virus to generate antibodies. The protein sequence  
15 may be determined from the cDNA sequence. Amino acid sequences may be analyzed by methods well known to those skilled in the art to determine whether they produce hydrophobic or hydrophilic regions in the proteins which they build. In the case of cell membrane proteins, hydrophobic regions are well known to form the part of the protein that is inserted into the lipid bilayer of the cell membrane, while hydrophilic regions are located  
20 on the cell surface, in an aqueous environment. Usually, the hydrophilic regions will be more immunogenic than the hydrophobic regions. Therefore the hydrophilic amino acid sequences may be selected and used to generate antibodies specific to polypeptide encoded by the isolated nucleic acid molecule encoding the DNA virus. The selected peptides may be prepared using commercially available machines. As an alternative, DNA, such as a  
25 cDNA or a fragment thereof, may be cloned and expressed and the resulting polypeptide recovered and used as an immunogen.

Polyclonal antibodies against these peptides may be produced by immunizing animals using the selected peptides. Monoclonal antibodies are prepared using hybridoma technology by  
30 fusing antibody producing B cells from immunized animals with myeloma cells and selecting the resulting hybridoma cell line producing the desired antibody. Alternatively, monoclonal antibodies may be produced by in vitro techniques known to a person of ordinary skill in the art. These antibodies are useful to detect the expression of polypeptide encoded by the isolated DNA molecule of the DNA virus in living animals, in humans, or in biological  
35 tissues or fluids isolated from animals or humans.

5

The antibodies may be detectably labelled, utilizing conventional labelling techniques well-known to the art. Thus, the antibodies may be radiolabelled using, for example, radioactive isotopes such as  $^3\text{H}$ ,  $^{125}\text{I}$ ,  $^{131}\text{I}$ , and  $^{35}\text{S}$ . The antibodies may also be labelled using fluorescent labels, enzyme labels, free radical labels, or bacteriophage labels, using techniques known in the art. Typical fluorescent labels include fluorescein isothiocyanate, rhodamine, phycoerythrin, phycocyanin, allophycocyanin, and Texas Red.

15 Since specific enzymes may be coupled to other molecules by covalent links, the possibility also exists that they might be used as labels for the production of tracer materials. Suitable enzymes include alkaline phosphatase, beta-galactosidase, glucose-6-phosphate dehydrogenase, maleate dehydrogenase, and peroxidase. Two principal types of enzyme immunoassay are the enzyme-linked immunosorbent assay (ELISA), and the homogeneous enzyme immunoassay, also known as enzyme-multiplied immunoassay (EMIT, Syva Corporation, Palo Alto, CA). In the ELISA system, separation may be achieved, for example, by the use of antibodies coupled to a solid phase. The EMIT system depends on  
20 deactivation of the enzyme in the tracer-antibody complex; the activity can thus be measured without the need for a separation step.

25 Additionally, chemiluminescent compounds may be used as labels. Typical chemiluminescent compounds include luminol, isoluminol, aromatic acridinium esters, imidazoles, acridinium salts, and oxalate esters. Similarly, bioluminescent compounds may be utilized for labelling, the bioluminescent compounds including luciferin, luciferase, and aequorin. Once labeled, the antibody may be employed to identify and quantify immunologic counterparts (antibody or antigenic polypeptide) utilizing techniques well-known to the art.

30

A description of a radioimmunoassay (RIA) may be found in *Laboratory Techniques in Biochemistry and Molecular Biology* [52], with particular reference to the chapter entitled "An Introduction to Radioimmune Assay and Related Techniques" by Chard, T., incorporated by reference herein. A description of general immunometric assays of various  
35 types can be found in the following U.S. Pat. Nos. 4,376,110 (David *et al.*) or 4,098,876

5 (Piasio).

One can use immunoassays to detect for the p-Hyde gene, specific peptides, or for antibodies to the virus or peptides. A general overview of the applicable technology is in Harlow and Lane [32], incorporated by reference herein.

10

In one embodiment, antibodies to the human p-Hyde can be used to detect the agent in the sample. In brief, to produce antibodies to the agent or peptides, the sequence being targeted is expressed in transfected cells, preferably bacterial cells, and purified. The product is injected into a mammal capable of producing antibodies. Either monoclonal or polyclonal  
15 antibodies (as well as any recombinant antibodies) specific for the gene product can be used in various immunoassays. Such assays include competitive immunoassays, radioimmunoassays, Western blots, ELISA, indirect immunofluorescent assays and the like. For competitive immunoassays, see Harlow and Lane [32] at pages 567-573 and 584-589.

20

In a further embodiment of this invention, commercial test kits suitable for use by a medical specialist may be prepared to determine the presence or absence of predetermined binding activity or predetermined binding activity capability to suspected target cells. In accordance with the testing techniques discussed above, one class of such kits will contain at least the labeled polypeptide or its binding partner, for instance an antibody specific thereto, and  
25 directions, of course, depending upon the method selected, e.g., "competitive," "sandwich," "DASP" and the like. The kits may also contain peripheral reagents such as buffers, stabilizers, etc.

30

Monoclonal antibodies or recombinant antibodies may be obtained by various techniques familiar to those skilled in the art. Briefly, spleen cells or other lymphocytes from an animal immunized with a desired antigen are immortalized, commonly by fusion with a myeloma cell (see, Kohler and Milstein [50], incorporated herein by reference). Alternative methods of immortalization include transformation with Epstein Barr Virus, oncogenes, or retroviruses, or other methods well known in the art. Colonies arising from single  
35 immortalized cells are screened for production of antibodies of the desired specificity and



5 affinity for the antigen, and yield of the monoclonal antibodies produced by such cells may be enhanced by various techniques, including injection into the peritoneal cavity of a vertebrate host. New techniques using recombinant phage antibody expression systems can also be used to generate monoclonal antibodies. See for example: McCafferty, J *et al.* [64]; Hoogenboom, H.R. *et al.* [39]; and Marks, J.D. *et al.* [60].

10

Such peptides may be produced by expressing the specific sequence in a recombinantly engineered cell such as bacteria, yeast, filamentous fungal, insect (especially employing baculoviral vectors), and mammalian cells. Those of skill in the art are knowledgeable in the numerous expression systems available for expression of herpes virus protein.

15

Briefly, the expression of natural or synthetic nucleic acids encoding viral protein will typically be achieved by operably linking the desired sequence or portion thereof to a promoter (which is either constitutive or inducible), and incorporated into an expression vector. The vectors are suitable for replication or integration in either prokaryotes or eukaryotes. Typical cloning vectors contain antibiotic resistance markers, genes for selection of transformants, inducible or regulatable promoter regions, and translation terminators that are useful for the expression of viral genes.

20

Methods for the expression of cloned genes in bacteria are also well known. In general, to obtain high level expression of a cloned gene in a prokaryotic system, it is advisable to construct expression vectors containing a strong promoter to direct mRNA transcription. The inclusion of selection markers in DNA vectors transformed in *E. coli* is also useful. Examples of such markers include genes specifying resistance to antibiotics. See [81] *supra*, for details concerning selection markers and promoters for use in *E. coli*. Suitable eukaryote hosts may include plant cells, insect cells, mammalian cells, yeast, and filamentous fungi.

25

30

The peptides derived from the nucleic acids, peptide fragments are produced by recombinant technology may be purified by standard techniques well known to those of skill in the art. Recombinantly produced sequences can be directly expressed or expressed as a fusion protein. The protein is then purified by a combination of cell lysis (*e.g.*, sonication) and

35

5 affinity chromatography. For fusion products, subsequent digestion of the fusion protein with an appropriate proteolytic enzyme releases the desired peptide.

The proteins may be purified to substantial purity by standard techniques well known in the art, including selective precipitation with such substances as ammonium sulfate, column  
10 chromatography, immunopurification methods, and others. See, for instance, Scopes, R. [84], incorporated herein by reference.

This invention is directed to analogs of the isolated nucleic acid and polypeptide which  
15 comprise the amino acid sequence as set forth above. The analog may have an N-terminal methionine or an N-terminal polyhistidine optionally attached to the N or COOH terminus of the polypeptide which comprise the amino acid sequence.

In another embodiment, this invention contemplates peptide fragments of the polypeptide which result from proteolytic digestion products of the polypeptide. In another embodiment,  
20 the derivative of the polypeptide has one or more chemical moieties attached thereto. In another embodiment the chemical moiety is a water soluble polymer. In another embodiment the chemical moiety is polyethylene glycol. In another embodiment the chemical moiety is mon-, di-, tri- or tetrapegylated. In another embodiment the chemical moiety is N-terminal monopegylated.

25 A polypeptide "fragment," "portion" or "segment" is a stretch of amino acid residues of at least about five to seven contiguous amino acids, often at least about seven to nine contiguous amino acids, typically at least about nine to 13 contiguous amino acids and, most preferably, at least about 20 to 30 or more contiguous amino acids.

30 The polypeptides of the present invention, if soluble, may be coupled to a solid-phase support, e.g., nitrocellulose, nylon, column packing materials (e.g., Sepharose beads), magnetic beads, glass wool, plastic, metal, polymer gels, cells, or other substrates. Such supports may take the form, for example, of beads, wells, dipsticks, or membranes.

35

5 "Target region" refers to a region of the nucleic acid which is amplified and/or detected. The term "target sequence" refers to a sequence with which a probe or primer will form a stable hybrid under desired conditions.

10 Attachment of polyethylene glycol (PEG) to compounds is particularly useful because PEG has very low toxicity in mammals (Carpenter et al., 1971). For example, a PEG adduct of adenosine deaminase was approved in the United States for use in humans for the treatment of severe combined immunodeficiency syndrome. A second advantage afforded by the conjugation of PEG is that of effectively reducing the immunogenicity and antigenicity of heterologous compounds. For example, a PEG adduct of a human protein might be useful  
15 for the treatment of disease in other mammalian species without the risk of triggering a severe immune response. The compound of the present invention may be delivered in a microencapsulation device so as to reduce or prevent an host immune response against the compound or against cells which may produce the compound. The compound of the present invention may also be delivered microencapsulated in a membrane, such as a liposome.

20 Numerous activated forms of PEG suitable for direct reaction with proteins have been described. Useful PEG reagents for reaction with protein amino groups include active esters of carboxylic acid or carbonate derivatives, particularly those in which the leaving groups are N-hydroxysuccinimide, p-nitrophenol, imidazole or 1-hydroxy-2-nitrobenzene-4-sulfonate. PEG derivatives containing maleimido or haloacetyl groups are useful reagents  
25 for the modification of protein free sulfhydryl groups. Likewise, PEG reagents containing amino hydrazine or hydrazide groups are useful for reaction with aldehydes generated by periodate oxidation of carbohydrate groups in proteins.

30 In one embodiment, the amino acid residues of the polypeptide described herein are preferred to be in the "L" isomeric form. In another embodiment, the residues in the "D" isomeric form can be substituted for any L-amino acid residue, as long as the desired functional property of lectin activity is retained by the polypeptide. NH<sub>2</sub> refers to the free amino group present at the amino terminus of a polypeptide. COOH refers to the free carboxy group  
35 present at the carboxy terminus of a polypeptide. Abbreviations used herein are in keeping

5 with standard polypeptide nomenclature, *J. Biol. Chem.*, 243:3552-59 (1969).

It should be noted that all amino-acid residue sequences are represented herein by formulae whose left and right orientation is in the conventional direction of amino-terminus to carboxy-terminus. Furthermore, it should be noted that a dash at the beginning or end of an  
10 amino acid residue sequence indicates a peptide bond to a further sequence of one or more amino-acid residues.

Synthetic polypeptide, prepared using the well known techniques of solid phase, liquid phase, or peptide condensation techniques, or any combination thereof, can include natural  
15 and unnatural amino acids. Amino acids used for peptide synthesis may be standard Boc (N-amino protected N-t-butyloxycarbonyl) amino acid resin with the standard deprotecting, neutralization, coupling and wash protocols of the original solid phase procedure of Merrifield (1963, *J. Am. Chem. Soc.* 85:2149-2154), or the base-labile N-amino protected 9-fluorenylmethoxycarbonyl (Fmoc) amino acids first described by Carpino and Han (1972,  
20 *J. Org. Chem.* 37:3403-3409). Thus, polypeptide of the invention may comprise D-amino acids, a combination of D- and L-amino acids, and various "designer" amino acids (*e.g.*, methyl amino acids, C-methyl amino acids, and N-methyl amino acids, etc.) to convey special properties. Synthetic amino acids include ornithine for lysine, fluorophenylalanine for phenylalanine, and norleucine for leucine or isoleucine. Additionally, by assigning  
25 specific amino acids at specific coupling steps, alpha-helices, alpha turns, beta sheets, beta-turns, and cyclic peptides can be generated.

In one aspect of the invention, the peptides may comprise a special amino acid at the C-terminus which incorporates either a CO<sub>2</sub>H or CONH<sub>2</sub> side chain to simulate a free glycine  
30 or a glycine-amide group. Another way to consider this special residue would be as a D or L amino acid analog with a side chain consisting of the linker or bond to the bead. In one embodiment, the pseudo-free C-terminal residue may be of the D or the L optical configuration; in another embodiment, a racemic mixture of D and L-isomers may be used.

35 In an additional embodiment, pyroglutamate may be included as the N-terminal residue of

5 the peptide. Although pyroglutamate is not amenable to sequence by Edman degradation,  
by limiting substitution to only 50% of the peptides on a given bead with N-terminal  
pyroglutamate, there will remain enough non-pyroglutamate peptide on the bead for  
sequencing. One of ordinary skill would readily recognize that this technique could be used  
for sequencing of any peptide that incorporates a residue resistant to Edman degradation at  
10 the N-terminus. Other methods to characterize individual peptides that demonstrate desired  
activity are described in detail *infra*. Specific activity of a peptide that comprises a blocked  
N-terminal group, *e.g.*, pyroglutamate, when the particular N-terminal group is present in  
50% of the peptides, would readily be demonstrated by comparing activity of a completely  
(100%) blocked peptide with a non-blocked (0%) peptide.

15 In addition, the present invention envisions preparing peptides that have more well defined  
structural properties, and the use of peptidomimetics, and peptidomimetic bonds, such as  
ester bonds, to prepare peptides with novel properties. In another embodiment, a peptide  
may be generated that incorporates a reduced peptide bond, *i.e.*,  $R_1-CH_2-NH-R_2$ , where  $R_1$   
20 and  $R_2$  are amino acid residues or sequences. A reduced peptide bond may be introduced as  
a dipeptide subunit. Such a molecule would be resistant to peptide bond hydrolysis, *e.g.*,  
protease activity. Such peptides would provide ligands with unique function and activity,  
such as extended half-lives *in vivo* due to resistance to metabolic breakdown, or protease  
activity. Furthermore, it is well known that in certain systems constrained peptides show  
25 enhanced functional activity (Hruby, 1982, Life Sciences 31:189-199; Hruby et al., 1990,  
Biochem J. 268:249-262); the present invention provides a method to produce a constrained  
peptide that incorporates random sequences at all other positions.



5 A constrained, cyclic or rigidized peptide may be prepared synthetically, provided that in at least two positions in the sequence of the peptide an amino acid or amino acid analog is inserted that provides a chemical functional group capable of cross-linking to constrain, cyclise or rigidize the peptide after treatment to form the cross-link. Cyclization will be favored when a turn-inducing amino acid is incorporated. Examples of amino acids capable of cross-linking a peptide are cysteine to form disulfide, aspartic acid to form a lactone or a lactase, and a chelator such as carboxyl-glutamic acid (Gla) (Bachem) to chelate a transition metal and form a cross-link. Protected carboxyl glutamic acid may be prepared by modifying the synthesis described by Zee-Cheng and Olson (1980, Biophys. Biochem. Res. Commun. 94:1128-1132). A peptide in which the peptide sequence comprises at least two amino acids capable of cross-linking may be treated, *e.g.*, by oxidation of cysteine residues to form a disulfide or addition of a metal ion to form a chelate, so as to cross-link the peptide and form a constrained, cyclic or rigidized peptide.

20 The present invention provides strategies to systematically prepare cross-links. For example, if four cysteine residues are incorporated in the peptide sequence, different protecting groups may be used (Hiskey, 1981, in *The Peptides: Analysis, Synthesis, Biology*, Vol. 3, Gross and Meienhofer, eds., Academic Press: New York, pp. 137-167; Ponsanti et al., 1990, *Tetrahedron* 46:8255-8266). The first pair of cysteine may be deprotected and oxidized, then the second set may be deprotected and oxidized. In this way a defined set of disulfide cross-links may be formed. Alternatively, a pair of cysteine and a pair of collating amino acid analogs may be incorporated so that the cross-links are of a different chemical nature.

The following non-classical amino acids may be incorporated in the peptide in order to introduce particular conformational motifs: 1,2,3,4-tetrahydroisoquinoline-3-carboxylate (Kazmierski et al., 1991, J. Am. Chem. Soc. 113:2275-2283); (2S,3S)-methyl-phenylalanine, (2S,3R)-methyl-phenylalanine, (2R,3S)-methyl-phenylalanine and (2R,3R)-methyl-phenylalanine (Kazmierski and Hruby, 1991, Tetrahedron Lett.); 2-aminotetrahydronaphthalene-2-carboxylic acid (Landis, 1989, Ph.D. Thesis, University of Arizona); hydroxy-1,2,3,4-tetrahydroisoquinoline-3-carboxylate (Miyake et al., 1989, J. Takeda Res. Labs. 43:53-76); ?-carboline (D and L) (Kazmierski, 1988, Ph.D. Thesis, University of Arizona); HIC (histidine isoquinoline carboxylic acid) (Zechel et al., 1991, Int. J. Pep. Protein Res. 43); and HIC (histidine cyclic urea) (Dharanipragada).

The following amino acid analogs and peptidomimetics may be incorporated into a peptide to induce or favor specific secondary structures: LL-Acp (LL-3-amino-2-propenidone-6-carboxylic acid), a dipeptide analog (Kemp et al., 1985, J. Org. Chem. 50:5834-5838); and analogs provided by the following references: Nagai and Sato, 1985, Tetrahedron Lett. 26:647-650; DiMaio et al., 1989, J. Chem. Soc. Perkin Trans. p. 1687; also a Gly-Ala turn analog (Kahn et al., 1989, Tetrahedron Lett. 30:2317); amide bond isostere (Jones et al., 1988, Tetrahedron Lett. 29:3853-3856); tetrazol (Zabrocki et al., 1988, J. Am. Chem. Soc. 110:5875-5880); DTC (Samanen et al., 1990, Int. J. Protein Pep. Res. 35:501:509); and analogs taught in Olson et al., 1990, J. Am. Chem. Sci. 112:323-333 and Garvey et al., 1990, J. Org. Chem. 56:436. Conformationally restricted mimetics of beta turns and beta bulges, and peptides containing them, are described in U.S. Patent No. 5,440,013, issued August 8, 1995 to Kahn.

The present invention further provides for modification or derivatization of the polypeptide or peptide of the invention. Modifications of peptides are well known to one of ordinary skill, and include phosphorylation, carboxymethylation, and acylation. Modifications may be effected by chemical or enzymatic means. In another aspect, glycosylated or fatty acylated peptide derivatives may be prepared. Preparation of glycosylated or fatty acylated peptides is well known in the art. Fatty acyl peptide derivatives may also be prepared. For example,

5 and not by way of limitation, a free amino group (N-terminal or lysyl) may be acylated, *e.g.*, myristoylated. In another embodiment an amino acid comprising an aliphatic side chain of the structure  $-(CH_2)_nCH_3$  may be incorporated in the peptide. This and other peptide-fatty acid conjugates suitable for use in the present invention are disclosed in U.K. Patent GB-8809162.4, International Patent Application PCT/AU89/00166, and reference 5, *supra*.

10

Mutations can be made in a nucleic acid encoding the polypeptide such that a particular codon is changed to a codon which codes for a different amino acid. Such a mutation is generally made by making the fewest nucleotide changes possible. A substitution mutation of this sort can be made to change an amino acid in the resulting protein in a non-conservative manner (i.e., by changing the codon from an amino acid belonging to a grouping of amino acids having a particular size or characteristic to an amino acid belonging to another grouping) or in a conservative manner (i.e., by changing the codon from an amino acid belonging to a grouping of amino acids having a particular size or characteristic to an amino acid belonging to the same grouping). Such a conservative change generally leads to less change in the structure and function of the resulting protein. A non-conservative change is more likely to alter the structure, activity or function of the resulting protein. The present invention should be considered to include sequences containing conservative changes which do not significantly alter the activity or binding characteristics of the resulting protein. Substitutes for an amino acid within the sequence may be selected from other members of the class to which the amino acid belongs. For example, the nonpolar (hydrophobic) amino acids include alanine, leucine, isoleucine, valine, proline, phenylalanine, tryptophan and methionine. Amino acids containing aromatic ring structures are phenylalanine, tryptophan, and tyrosine. The polar neutral amino acids include glycine, serine, threonine, cysteine, tyrosine, asparagine, and glutamine. The positively charged (basic) amino acids include arginine, lysine and histidine. The negatively charged (acidic) amino acids include aspartic acid and glutamic acid. Such alterations will not be expected to affect apparent molecular weight as determined by polyacrylamide gel electrophoresis, or isoelectric point.

20

25

30

35

Particularly preferred substitutions are:

- Lys for Arg and vice versa such that a positive charge may be maintained;

- 5       - Glu for Asp and vice versa such that a negative charge may be maintained;  
      - Ser for Thr such that a free -OH can be maintained; and  
      - Gln for Asn such that a free NH<sub>2</sub> can be maintained.

10       Synthetic DNA sequences allow convenient construction of genes which will express analogs  
      or "muteins". A general method for site-specific incorporation of unnatural amino acids into  
      proteins is described in Noren, et al. *Science*, 244:182-188 (April 1989). This method may  
      be used to create analogs with unnatural amino acids.

15       In accordance with the present invention there may be employed conventional molecular  
      biology, microbiology, and recombinant DNA techniques within the skill of the art. Such  
      techniques are explained fully in the literature. See, e.g., Sambrook et al, "Molecular  
      Cloning: A Laboratory Manual" (1989); "Current Protocols in Molecular Biology" Volumes  
      I-III [Ausubel, R. M., ed. (1994)]; "Cell Biology: A Laboratory Handbook" Volumes I-III [J.  
      E. Celis, ed. (1994)]; "Current Protocols in Immunology" Volumes I-III [Coligan, J. E., ed.  
20       (1994)]; "Oligonucleotide Synthesis" (M.J. Gait ed. 1984); "Nucleic Acid Hybridization"  
      [B.D. Hames & S.J. Higgins eds. (1985)]; "Transcription And Translation" [B.D. Hames &  
      S.J. Higgins, eds. (1984)]; "Animal Cell Culture" [R.I. Freshney, ed. (1986)]; "Immobilized  
      Cells And Enzymes" [IRL Press, (1986)]; B. Perbal, "A Practical Guide To Molecular  
      Cloning" (1984).

25       In an additional embodiment, pyroglutamate may be included as the N-terminal residue of  
      the peptide. Although pyroglutamate is not amenable to sequence by Edman degradation,  
      by limiting substitution to only 50% of the peptides on a given bead with N-terminal  
      pyroglutamate, there will remain enough non-pyroglutamate peptide on the bead for  
30       sequencing. One of ordinary skill in would readily recognize that this technique could be  
      used for sequencing of any peptide that incorporates a residue resistant to Edman degradation  
      at the N-terminus. Other methods to characterize individual peptides that demonstrate  
      desired activity are described in detail *infra*. Specific activity of a peptide that comprises a  
      blocked N-terminal group, e.g., pyroglutamate, when the particular N-terminal group is  
35       present in 50% of the peptides, would readily be demonstrated by comparing activity of a

5 completely (100%) blocked peptide with a non-blocked (0%) peptide.

*Chemical Moieties For Derivatization.* Chemical moieties suitable for derivatization may be selected from among water soluble polymers. The polymer selected should be water soluble so that the component to which it is attached does not precipitate in an aqueous  
10 environment, such as a physiological environment. Preferably, for therapeutic use of the end-product preparation, the polymer will be pharmaceutically acceptable. One skilled in the art will be able to select the desired polymer based on such considerations as whether the polymer/component conjugate will be used therapeutically, and if so, the desired dosage, circulation time, resistance to proteolysis, and other considerations. For the present  
15 component or components, these may be ascertained using the assays provided herein.

The water soluble polymer may be selected from the group consisting of, for example, polyethylene glycol, copolymers of ethylene glycol/propylene glycol, carboxymethylcellulose, dextran, polyvinyl alcohol, polyvinyl pyrrolidone, poly-1,  
20 3-dioxolane, poly-1,3,6-trioxane, ethylene/maleic anhydride copolymer, polyaminoacids (either homopolymers or random copolymers), and dextran or poly(n-vinyl pyrrolidone)polyethylene glycol, propylene glycol homopolymers, polypropylene oxide/ethylene oxide co- polymers, polyoxyethylated polyols and polyvinyl alcohol. Polyethylene glycol propionaldehyde may have advantages in manufacturing due to its  
25 stability in water.

The number of polymer molecules so attached may vary, and one skilled in the art will be able to ascertain the effect on function. One may mono-derivatize, or may provide for a di-,  
30 tri-, tetra- or some combination of derivatization, with the same or different chemical moieties (e.g., polymers, such as different weights of polyethylene glycols). The proportion of polymer molecules to component or components molecules will vary, as will their concentrations in the reaction mixture. In general, the optimum ratio (in terms of efficiency of reaction in that there is no excess unreacted component or components and polymer) will  
35 be determined by factors such as the desired degree of derivatization (e.g., mono, di-, tri-,



5 etc.), the molecular weight of the polymer selected, whether the polymer is branched or unbranched, and the reaction conditions.

10 This invention provides a method for determining whether a subject carries a mutation in the p-Hyde gene which comprises: a) obtaining an appropriate nucleic acid sample from the subject; and (b) determining whether the nucleic acid sample from step (a) is, or is derived from, a nucleic acid which encodes mutant p-Hyde so as to thereby determine whether a subject carries a mutation in the p-Hyde gene. In one embodiment, the nucleic acid sample in step (a) comprises mRNA corresponding to the transcript of DNA encoding a mutant p-Hyde, and wherein the determining of step (b) comprises: (i) contacting the mRNA with the  
15 oligonucleotide under conditions permitting binding of the mRNA to the oligonucleotide so as to form a complex; (ii) isolating the complex so formed; and (iii) identifying the mRNA in the isolated complex so as to thereby determine whether the mRNA is, or is derived from, a nucleic acid which encodes mutant p-Hyde. In another embodiment, the determining of step (b) comprises: i) contacting the nucleic acid sample of step (a), and the isolated nucleic acid  
20 with restriction enzymes under conditions permitting the digestion of the nucleic acid sample, and the isolated nucleic acid into distinct, distinguishable pieces of nucleic acid; (ii) isolating the pieces of nucleic acid; and (iii) comparing the pieces of nucleic acid derived from the nucleic acid sample with the pieces of nucleic acid derived from the isolated nucleic acid so as to thereby determine whether the nucleic acid sample is, or is derived from, a  
25 nucleic acid which encodes mutant p-Hyde.

The present invention further provides methods of preparing a polynucleotide comprising polymerizing nucleotides to yield a sequence comprised of at least eight consecutive nucleotides of the p-Hyde gene; and methods of preparing a polypeptide  
30 comprising polymerizing amino acids to yield a sequence comprising at least five amino acids encoded within the p-Hyde gene.

This invention provides a pharmaceutical composition comprising an amount of the polypeptide and a pharmaceutically effective carrier or diluent.

35

5 This invention provides a method for determining whether a subject carries a mutation in  
the p-Hyde gene which comprises: (a) obtaining an appropriate nucleic acid sample from the  
subject; and (b) determining whether the nucleic acid sample from step (a) is, or is derived  
from, a nucleic acid which encodes mutant p-Hyde so as to thereby determine whether a  
subject carries a mutation in the p-Hyde gene. In one embodiment the nucleic acid sample  
10 in step (a) comprises mRNA corresponding to the transcript of DNA encoding a mutant p-  
Hyde, and wherein the determining of step (b) comprises: (i) contacting the mRNA with the  
oligonucleotide under conditions permitting binding of the mRNA to the oligonucleotide so  
as to form a complex; (ii) isolating the complex so formed; and (iii) identifying the mRNA  
in the isolated complex so as to thereby determine whether the mRNA is, or is derived from,  
15 a nucleic acid which encodes mutant p-Hyde. In another embodiment the determining of step  
(b) comprises: (i) contacting the nucleic acid sample of step (a), and the isolated nucleic acid  
of claim 1 with restriction enzymes under conditions permitting the digestion of the nucleic  
acid sample, and the isolated nucleic acid into distinct, distinguishable pieces of nucleic acid;  
(ii) isolating the pieces of nucleic acid; and (iii) comparing the pieces of nucleic acid derived  
20 from the nucleic acid sample with the pieces of nucleic acid derived from the isolated nucleic  
acid so as to thereby determine whether the nucleic acid sample is, or is derived from, a  
nucleic acid which encodes mutant p-Hyde.

Detection of point mutations or variations may be accomplished by molecular cloning of the  
25 p-Hyde allele(s) and sequencing the allele(s) using techniques well known in the art.  
Alternatively, the gene sequences can be amplified directly from a genomic DNA  
preparation from the tumor tissue, using known techniques. The DNA sequence of the  
amplified sequences can then be determined. There are six well known methods for a  
more complete, yet still indirect, test for confirming the presence of a susceptibility allele:  
30 1) single stranded conformation analysis (SSCA) (Orita et al., 1989); 2) denaturing gradient  
gel electrophoresis (DGGE) (Wartell et al., 1990; Sheffield et al., 1989); 3) RNase  
protection assays (Finkelstein et al., 1990; Kinszler et al., 1991); 4) allele-specific  
oligonucleotides (ASOs) (Conner et al., 1983); 5) the use of proteins which recognize  
nucleotide mismatches, such as the E. coli mutS protein (Modrich, 1991); and 6)  
35 allele-specific PCR (Rano & Kidd, 1989). For allele-specific PCR, primers are used

5 which hybridize at their 3' ends to a particular p-Hyde mutation. If the particular p-Hyde mutation is not present, an amplification product is not observed. Amplification Refractory Mutation System (ARMS) can also be used, is disclosed in European Patent Application Publication No. 0332435 and in Newton et al., 1989. Insertions and deletions of genes can also be detected by cloning, sequencing and amplification. In addition,  
10 restriction fragment length polymorphism (RFLP) probes for the gene or surrounding marker genes can be used to score alteration of an allele or an insertion in a polymorphic fragment. Such a method is particularly useful for screening relatives of an affected individual for the presence of the p-Hyde mutation found in that individual. Other techniques for detecting insertions and deletions as known in the art can be used.

15 In similar fashion, DNA probes can be used to detect mismatches, through enzymatic or chemical cleavage. See, e.g., Cotton et al., 1988; Shenk et al., 1975; Novack et al., 1986. Alternatively, mismatches can be detected by shifts in the electrophoretic mobility of mismatched duplexes relative to matched duplexes. See, e.g., Cariello, 1988. With either  
20 riboprobes or DNA probes, the cellular mRNA or DNA which might contain a mutation can be amplified using PCR (see below) before hybridization. Changes in DNA of the p-Hyde gene can also be detected using Southern hybridization, especially if the changes are gross rearrangements, such as deletions and insertions.

25 As noted above, non-PCR based screening assays are also contemplated in this invention. This procedure hybridizes a nucleic acid probe (or an analog such as a methyl phosphonate backbone replacing the normal phosphodiester), to the low level DNA target. This probe may have an enzyme covalently linked to the probe, such that the covalent linkage does not interfere with the specificity of the hybridization. This  
30 enzyme-probe-conjugate-target nucleic acid complex can then be isolated away from the free probe enzyme conjugate and a substrate is added for enzyme detection. Enzymatic activity is observed as a change in color development or luminescent output resulting in a  $10^3$  -  $10^6$  increase in sensitivity. For an example relating to the preparation of oligodeoxynucleotide-alkaline phosphatase conjugates and their use as hybridization  
35 probes see Jablonski et al., 1986.

5 Two-step label amplification methodologies are known in the art. These assays work on  
the principle that a small ligand (such as digoxigenin, biotin, or the like) is attached to  
a nucleic acid probe capable of specifically binding p-Hyde. Allele specific probes are also  
contemplated within the scope of this example. In one example, the small ligand attached  
to the nucleic acid probe is specifically recognized by an antibody-enzyme conjugate. In one  
10 embodiment of this example, digoxigenin is attached to the nucleic acid probe.  
Hybridization is detected by an antibody-alkaline phosphatase conjugate which turns over  
a chemiluminescent substrate. For methods for labeling nucleic acid probes according to  
this embodiment see Martin et al. , 1990.

15 In a second example, the small ligand is recognized by a second ligand-enzyme  
conjugate that is capable of specifically complexing to the first ligand. A well known  
embodiment of this example is the biotin-avidin type of interactions. For methods for  
labeling nucleic acid probes and their use in biotin-avidin based assays see Rigby et al.,  
1977 and Nguyen et al., 1992.

20 It is also contemplated within the scope of this invention that the nucleic acid probe  
assays of this invention will employ a cocktail of nucleic acid probes capable of  
detecting p-Hyde. Thus, in one example to detect the presence of p-Hyde in a cell  
sample, more than one probe complementary to p-Hyde is employed and in particular  
25 the number of different probes is alternatively 2, 3, or 5 different nucleic acid probe  
sequences. In another example, to detect the presence of mutations in the p-Hyde gene  
sequence in a patient, more than one probe complementary to p-Hyde is employed where  
the cocktail includes probes capable of patients with alternate-specific mutations identified  
in populations of patients with alterations in p-Hyde. In this embodiment, any number of  
30 probes can be used, and will preferably include probes corresponding to the major gene  
mutations identified as predisposing an individual to breast cancer.

This invention provides a method for screening a tumor sample from a human subject for a  
somatic alteration in a p-Hyde gene in said tumor which comprises gene comparing a first  
35 sequence selected from the group consisting of a p-Hyde gene from said tumor sample, p-

5 Hyde RNA from said tumor sample and p-Hyde cDNA made from mRNA from said tumor  
sample with a second sequence selected from the group consisting of p-Hyde gene from a  
nontumor sample of said subject, p-Hyde RNA from said nontumor sample and p-Hyde  
cDNA made from mRNA from said nontumor sample, wherein a difference in the sequence  
10 of the p-Hyde gene, p-Hyde RNA or p-Hyde cDNA from said tumor sample from the  
sequence of the p-Hyde gene, p-Hyde RNA or p-Hyde cDNA from said nontumor sample  
indicates a somatic alteration in the p-Hyde gene in said tumor sample.

This invention provides a method for screening a tumor sample from a human subject for the  
presence of a somatic alteration in a p-Hyde gene in said tumor which comprises comparing  
15 p-Hyde polypeptide from said tumor sample from said subject to p-Hyde polypeptide from  
a nontumor sample from said subject to analyze for a difference between the polypeptides,  
wherein said comparing is performed by (i) detecting either a full length polypeptide or a  
truncated polypeptide in each sample or (ii) contacting an antibody which specifically binds  
to either an epitope of an altered p-Hyde polypeptide or an epitope of a wild-type p-Hyde  
20 polypeptide to the p-Hyde polypeptide from each sample and detecting antibody binding,  
wherein a difference between the p-Hyde polypeptide from said tumor sample from the p-  
Hyde polypeptide from said nontumor sample indicates the presence of a somatic alteration  
in the p-Hyde gene in said tumor sample.

This invention is particularly useful for screening compounds by using the p-Hyde  
25 polypeptide or binding fragment thereof in any of a variety of drug screening techniques.  
The p-Hyde polypeptide or fragment employed in such a test may either be free in solution,  
affixed to a solid support, or borne on a cell surface. One method of drug screening utilizes  
eucaryotic or procaryotic host cells which are stably transformed with recombinant  
polynucleotides expressing the polypeptide or fragment, preferably in competitive binding  
30 assays. Such cells, either in viable or fixed form, can be used for standard binding assays.  
One may measure, for example, for the formation of complexes between a p-Hyde  
polypeptide or fragment and the agent being tested, or examine the degree to which the  
formation of a complex between a p-Hyde polypeptide or fragment and a known ligand is  
interfered with by the agent being tested.

35



5 Thus, the present invention provides methods of screening for drugs comprising  
contacting such an agent with a p-Hyde polypeptide or fragment thereof and assaying (i)  
for the presence of a complex between the agent and the p-Hyde polypeptide or fragment,  
or (ii) for the presence of a complex between the p-Hyde polypeptide or fragment and a  
ligand, by methods well known in the art. In such competitive binding assays the p-  
10 Hyde polypeptide or fragment is typically labeled. Free p-Hyde polypeptide or fragment  
is separated from that present in a protein:protein complex, and the amount of free (i.e.,  
uncomplexed) label is a measure of the binding of the agent being tested to p-Hyde or its  
interference with p-Hyde:ligand binding, respectively.

15 Another technique for drug screening provides high throughput screening for compounds  
having suitable binding affinity to the p-Hyde polypeptides and is described in detail in  
Geysen, PCT published application WO 84/03564, published on Sep. 13, 1984. Briefly  
stated, large numbers of different small peptide test compounds are synthesized on a  
solid substrate, such as plastic pins or some other surface. The peptide test compounds  
20 are reacted with p-Hyde polypeptide and washed. Bound p-Hyde polypeptide is then  
detected by methods well known in the art.

Purified p-Hyde can be coated directly onto plates for use in the aforementioned drug  
screening techniques. However, non-neutralizing antibodies to the polypeptide can be  
25 used to capture antibodies to immobilize the p-Hyde polypeptide on the solid phase. This  
invention also contemplates the use of competitive drug screening assays in which  
neutralizing antibodies capable of specifically binding the p-Hyde polypeptide compete  
with a test compound for binding to the p-Hyde polypeptide or fragments thereof. In this  
manner, the antibodies can be used to detect the presence of any peptide which shares  
30 one or more antigenic determinants of the p-Hyde polypeptide.

A further technique for drug screening involves the use of host eukaryotic cell lines  
or cells (such as described above) which have a nonfunctional p-Hyde gene. These host  
cell lines or cells are defective at the p-Hyde polypeptide level. The host cell lines or cells  
35 are grown in the presence of drug compound. The rate of growth of the host cells is measured

5 to determine if the compound is capable of regulating the growth of p-Hyde defective cells.

10 The goal of rational drug design is to produce structural analogs of biologically active polypeptides of interest or of small molecules with which they interact (e.g., agonists, antagonists, inhibitors) in order to fashion drugs which are, for example, more active or stable forms of the polypeptide, or which, e.g., enhance or interfere with the function of a polypeptide in vivo. See, e.g., Hodgson, 1991. In one approach, one first determines the three-dimensional structure of a protein of interest (e.g., p-Hyde polypeptide) or, for example, of the p-Hyde-receptor or ligand complex, by x-ray crystallography, by computer modeling or most typically, by a combination of approaches. Less often, useful information  
15 regarding the structure of a polypeptide may be gained by modeling based on the structure of homologous proteins. An example of rational drug design is the development of HIV protease inhibitors (Erickson et al., 1990). In addition, peptides (e.g., p-Hyde polypeptide) are analyzed by an alanine scan (Wells, 1991). In this technique, an amino acid residue is replaced by Ala, and its effect on the peptide's activity is determined. Each of the  
20 amino acid residues of the peptide is analyzed in this manner to determine the important regions of the peptide.

It is also possible to isolate a target-specific antibody, selected by a functional assay, and then to solve its crystal structure. In principle, this approach yields a pharmacore upon which  
25 subsequent drug design can be based. It is possible to bypass protein crystallography altogether by generating anti-idiotypic antibodies (anti-ids) to a functional, pharmacologically active antibody. As a mirror image of a mirror image, the binding site of the anti-ids would be expected to be an analog of the original receptor. The anti-id could then be used to identify and isolate peptides from banks of chemically or  
30 biologically produced banks of peptides. Selected peptides would then act as the pharmacore.

Thus, one may design drugs which have, e.g., improved p-Hyde polypeptide activity or stability or which act as inhibitors, agonists, antagonists, etc. of p-Hyde polypeptide activity.  
35 By virtue of the availability of cloned p-Hyde sequences, sufficient amounts of the p-Hyde

5 polypeptide may be made available to perform such analytical studies as x-ray crystallography. In addition, the knowledge of the p-Hyde protein sequence provided herein will guide those employing computer modeling techniques in place of, or in addition to x-ray crystallography.

10 This invention provides a method for identifying a chemical compound which is capable inducing susceptibility to cell death which comprises: (a) contacting the p-Hyde with a chemical compound under conditions permitting binding between the p-Hyde and the chemical compound; (b) detecting specific binding of the chemical compound to the p-Hyde; and (c) determining whether the chemical compound inhibits the p-Hyde so as to identify a  
15 chemical compound which is capable of capable inducing susceptibility to cell death. Useful diagnostic techniques include, but are not limited to fluorescent in situ hybridization (FISH), direct DNA sequencing, PFGE analysis, Southern blot analysis, single stranded conformation analysis (SSCA), Rnase protection assay, allele-specific oligonucleotide (ASO), dot blot analysis and PCR-SSCP, as discussed in detail further below.

20 A rapid preliminary analysis to detect polymorphisms in DNA sequences can be performed by looking at a series of Southern blots of DNA cut with one or more restriction enzymes, preferably with a large number of restriction enzymes. Each blot contains a series of normal individuals and a series of tumors. Southern blots displaying hybridizing fragments  
25 (differing in length from control DNA when probed with sequences near or including the p-Hyde gene) indicate a possible mutation. If restriction enzymes which produce very large restriction fragments are used, then pulsed field gel electrophoresis (PFGE) is employed.

30 Detection of point mutations may be accomplished by molecular cloning of the p-Hyde allele(s) and sequencing the allele(s) using techniques well known in the art. Alternatively, the gene sequences can be amplified directly from a genomic DNA preparation from the tumor tissue, using known techniques. The DNA sequence of the amplified sequences can then be determined. There are six well known methods for a more  
35 complete, yet still indirect, test for confirming the presence of a susceptibility allele: 1)

5 single stranded conformation analysis (SSCA) (Orita et al., 1989); 2) denaturing gradient  
gel electrophoresis (DGGE) (Wartell et al., 1990; Sheffield et al., 1989); 3) RNase  
protection assays (Finkelstein et al., 1990; Kinszler et al., 1991); 4) allele-specific  
oligonucleotides (ASOs) (Conner et al., 1983); 5) the use of proteins which recognize  
10 nucleotide mismatches, such as the *E. coli* mutS protein (Modrich, 1991); and 6)  
allele-specific PCR (Rano & Kidd, 1989). For allele-specific PCR, primers are used  
which hybridize at their 3' ends to a particular p-Hyde mutation. If the particular p-  
Hyde mutation is not present, an amplification product is not observed. Amplification  
Refractory Mutation System (ARMS) can also be used, as disclosed in European Patent  
Application Publication No. 0332435 and in Newton et al., 1989. Insertions and deletions  
15 of genes can also be detected by cloning, sequencing and amplification. In addition,  
restriction fragment length polymorphism (RFLP) probes for the gene or surrounding marker  
genes can be used to score alteration of an allele or an insertion in a polymorphic  
fragment. Such a method is particularly useful for screening relatives of an affected  
individual for the presence of the p-Hyde mutation found in that individual. Other  
20 techniques for detecting insertions and deletions as known in the art can be used.

In similar fashion, DNA probes can be used to detect mismatches, through enzymatic or  
chemical cleavage. See, e.g., Cotton et al., 1988; Shenk et al., 1975; Novack et al., 1986.  
Alternatively, mismatches can be detected by shifts in the electrophoretic mobility of  
25 mismatched duplexes relative to matched duplexes. See, e.g., Cariello, 1988. With either  
riboprobes or DNA probes, the cellular mRNA or DNA which might contain a mutation can  
be amplified using PCR (see below) before hybridization. Changes in DNA of the p-Hyde  
gene can also be detected using Southern hybridization, especially if the changes are gross  
rearrangements, such as deletions and insertions.

30 DNA sequences of the p-Hyde gene which have been amplified by use of PCR may also be  
screened using allele-specific probes. These probes are nucleic acid oligomers, each of  
which contains a region of the p-Hyde gene sequence harboring a known mutation. For  
example, one oligomer may be about 30 nucleotides in length, corresponding to a  
35 portion of the p-Hyde gene sequence. By use of a battery of such allele-specific

5 probes, PCR amplification products can be screened to identify the presence of a previously identified mutation in the p-Hyde gene. Hybridization of allele-specific probes with amplified p-Hyde sequences can be performed, for example, on a nylon filter. Hybridization to a particular probe under stringent hybridization conditions indicates the presence of the same mutation in the tumor tissue as in the allele-specific probe.

10

Alteration of p-Hyde mRNA expression can be detected by any techniques known in the art. These include Northern blot analysis, PCR amplification and RNase protection. Diminished mRNA expression indicates an alteration of the wild-type p-Hyde gene. Alteration of wild-type p-Hyde genes can also be detected by screening for alteration of wild-type p-Hyde protein. For example, monoclonal antibodies immunoreactive with p-Hyde can be used to screen a tissue. Lack of cognate antigen would indicate a p-Hyde mutation. Antibodies specific for products of mutant alleles could also be used to detect mutant p-Hyde gene product. Such immunological assays can be done in any convenient formats known in the art. These include Western blots, immunohistochemical assays and ELISA assays. Any means for detecting an altered p-Hyde protein can be used to detect alteration of wild-type p-Hyde genes. Functional assays, such as protein binding determinations, can be used. In addition, assays can be used which detect p-Hyde biochemical function. Finding a mutant p-Hyde gene product indicates alteration of a wild-type p-Hyde gene. Mutant p-Hyde genes or gene products can also be detected in other human body samples, such as serum, stool, urine and sputum.

25

The present invention also provides for fusion polypeptides, comprising p-Hyde polypeptides and fragments. Homologous polypeptides may be fusions between two or more p-Hyde polypeptide sequences or between the sequences of p-Hyde and a related protein. Likewise, heterologous fusions may be constructed which would exhibit a combination of properties or activities of the derivative proteins. For example, ligand-binding or other domains may be "swapped" between different new fusion polypeptides or fragments. Such homologous or heterologous fusion polypeptides may display, for example, altered strength or specificity of binding. Fusion partners include immunoglobulins, bacterial beta -galactosidase, trpE, protein A, beta -lactamase, alpha

30

35



5        amylase, alcohol dehydrogenase and yeast alpha mating factor. See, e.g., Godowski et al.  
      , 1988. Fusion proteins will typically be made by either recombinant nucleic acid methods,  
      as described below, or may be chemically synthesized. Techniques for the synthesis of  
      polypeptides are described, for example, in Merrifield, 1963.

10       Probes for p-Hyde alleles may be derived from the sequences of the p-Hyde region or its  
      cDNAs. The probes may be of any suitable length, which span all or a portion of the p-  
      Hyde region, and which allow specific hybridization to the p-Hyde region. If the target  
      sequence contains a sequence identical to that of the probe, the probes may be short, e.g.,  
15       in the range of about 8-30 base pairs, since the hybrid will be relatively stable under even  
      stringent conditions. If some degree of mismatch is expected with the probe, i.e., if it  
      is suspected that the probe will hybridize to a variant region, a longer probe may be  
      employed which hybridizes to the target sequence with the requisite specificity.

20       The probes will include an isolated polynucleotide attached to a label or reporter molecule  
      and may be used to isolate other polynucleotide sequences, having sequence similarity  
      by standard methods. For techniques for preparing and labeling probes see, e.g.,  
      Sambrook et al., 1989 or Ausubel et al., 1992.

25       Probes comprising synthetic oligonucleotides or other polynucleotides of the present  
      invention may be derived from naturally occurring or recombinant single- or  
      double-stranded polynucleotides, or be chemically synthesized. Probes may also be labeled  
      by nick translation, Klenow fill-in reaction, or other methods known in the art.

30       Portions of the polynucleotide sequence having at least about eight nucleotides, usually  
      at least about 15 nucleotides, and fewer than about 6 kb, usually fewer than about 1.0 kb,  
      from a polynucleotide sequence encoding p-Hyde are preferred as probes. The probes may  
      also be used to determine whether mRNA encoding p-Hyde is present in a cell or tissue.

35       This invention provides a method of inhibiting the growth of cancer cells, comprising the  
      steps of obtaining the cells and contacting the cells of the subject with a replication-deficient  
      adenovirus type 5 expression vector comprising an adenovirus genome having a deletion in

5 the E1 and E3 region of the genome and an insertion within the region of a nucleic acid encoding p-Hyde under the control of a Rous Sarcoma virus promoter, thereby inhibiting the growth of the prostate cancer cells.

10 This invention provides a method of inhibiting the growth a prostate cancer cells, comprising: 1) obtaining a sample of prostate cells from a subject; 2) contacting the cells with a replication deficient adenovirus type 5 expression vector which comprises an adenovirus genome having a deletion in the E1 and E3 regions of the genome and an insertion within the regions of a p-Hyde cDNA under the control of a Rous Sarcoma virus promoter; and 3) introducing the cells into the subject, thereby inhibiting the growth of the cancer cells.

20 This invention provides a method of suppressing the growth of cancer cells in a subject, comprising introducing into the cancer cell an amount of a nucleic acid encoding a p-Hyde protein, a nucleic acid encoding a fragment of p-Hyde protein, or the nucleic acid encoding a mutant p-Hyde protein, thereby suppressing the growth of cancer cells in the subject.

25 This invention provides a method of suppressing growth of cancer cells in a subject, comprising administering to the subject a pharmaceutical composition comprising a therapeutically effective amount of a nucleic acid encoding a p-Hyde protein, a nucleic acid encoding a fragment of p-Hyde protein, or the nucleic acid encoding a mutant p-Hyde protein and a pharmaceutical acceptable carrier or diluent, thereby suppressing the growth of cancer cells in the subject.

30 As demonstrated herein apoptotic response was assessed using DNA laddering assay. DNAs were extracted after the respective treatment (24 hours with 1 mM Hydroxyurea followed by 24 hours with 0.1 mM 5'-dFURd) and analyzed on 1.6% agarose gel electrophoresis. In agreement with cell cycle analyses, apoptotic response of the stable transfectants AT1-H1 and AT3-H1 (pcHYDE transfected AT-1 and AT-3) are consistently and significantly higher relative to both parental (AT-1 and AT-3) and pcDNA-transfected parental cell lines (AT1-pc1 and AT3-pc1). In particular, the highest apoptotic response occurred in synchronized

5 culture under the induction with 0.1 mM 5'-dFUrd as shown in Fig.15. The enhanced  
apoptotic response in AT1-H1 and AT3-H1 transfectant after hydroxyurea treatment is the  
result of "thymineless death" (Kyprianou, 1994, Kyprianou et al., 1994) leading to depletion  
of intracellular thymidine-5-triphosphate (TTP) pools through indirect inhibition of  
thymidylate synthetase by fluorodeoxyuridine. However, the exact mechanism of the  
10 apoptosis itself in association with TTP depletion is not known. Taken together, these data  
suggest that the apoptotic response in the pcHYDE stable transfectants is likely due to the  
downstream effect of pcHYDE gene product.

15 This invention provides a method of inducing susceptibility to apoptosis of cancer cells,  
comprising introducing into the cancer cell an amount of a nucleic acid encoding a p-Hyde  
protein, a nucleic acid encoding a fragment of p-Hyde protein, or the nucleic acid encoding  
a mutant p-Hyde protein, thereby inducing susceptibility to apoptosis.

20 This invention provides a method of inducing susceptibility to apoptosis of cancer cells in  
a subject, comprising administering to the subject a pharmaceutical composition comprising  
a therapeutically effective amount of a nucleic acid encoding a p-Hyde protein, a nucleic  
acid encoding a fragment of p-Hyde protein, or the nucleic acid encoding a mutant p-Hyde  
protein and a pharmaceutical acceptable carrier or diluent, thereby inducing susceptibility to  
apoptosis.

25 As demonstrated herein, p-Hyde suppresses cancer growth *in vivo*. The lowest level of *p-Hyde*  
expression was observed in AT3 cell line as shown in Fig. 9. For this reason, AT3 cell  
line was transfected with pcHYDE, a construct of *p-Hyde* in mammalian expression vector  
of pcDNA3.1(-) under G418 selection. As negative control, AT-3 cell line was also  
30 transfected with the vector only and the stable transfectants obtained was designated as AT3-  
pc. The tumor growth of the parental cell line of AT-3 and in stable transfectant AT3-H1 and  
AT3-H2 have been evaluated *in vivo*. One million cells of each cell lines in 0.3 ml of Hanks  
solution were inoculated subcutaneously in each flank of inbred male Copenhagen rat. In  
this initial experiment, three groups of each five rats were injected with each cell line. The  
35 size of tumors were scored after a time schedule shown in Fig.18. These preliminary results

5 indicated that both AT3-H1 and AT3-H2 stable transfectant grew significantly slower than the AT-3 parental cell line demonstrating that the tumor growth regression in both stable transfectants and regulated by the pcHYDE gene product.

10 Interestingly, *p-Hyde* has the dual ability to act like a tumor suppressor gene and induce susceptibility to apoptosis by what may be p53 independent pathways. The growth of prostate tumors in rats was greatly inhibited by *p-Hyde*. Moreover, prostate cancer cells expressing *p-Hyde* were more sensitive to UV DNA damage driving these cells into cell programmed death. Analysis of DNA repair enzyme activity suggests a defect resulting in the presence of intact (6-4) PP and decreased cell survival by colony forming assay. However, the capacity of *p-Hyde* to induce susceptibility to apoptosis is not limited to UV DNA damage. 15 Chemotherapy agent, Fluorodeoxyuridine, a pyrimidine antimetabolite which is related to fluorouracil (5-FU) and has been used for treatment of a wide variety of human epithelial malignancies, also more readily induces apoptosis in prostate cancer cell expressing *p-Hyde*. Moreover, cancer cells expressing *p-Hyde* are also more susceptible to gamma radiation. 20 Thus, the mechanisms of cellular DNA injury are different for UV, gamma radiation, and Fluorodeoxyuridine suggesting that the ability to make cells more susceptible to apoptosis is more global in action. This unique function of *p-Hyde* may represent a new class genes that induce susceptibility to apoptosis. This is different than the function ascribed to tumor suppressor genes like p53 which directly induces apoptosis, not sensitivity to apoptosis 25 (Yonish-Rouach *et al.*, 1991). Moreover, *p-Hyde* activity is in contrast to *bcl-2* where the absence of *bcl-2*, not the presence of, makes the cancer cell more susceptible to cell programmed death (McDonnellet *et al.*, 1992).

30 This invention provides a method of suppressing cancer cells, comprising introducing into the cancer cell an amount of a nucleic acid encoding a p-Hyde protein, a nucleic acid encoding a fragment of p-Hyde protein, or the nucleic acid encoding a mutant p-Hyde protein, thereby inducing susceptibility to apoptosis.

35 This invention provides a method of suppressing cancer cells a subject, comprising administering to the subject a pharmaceutical composition comprising a therapeutically

5 effective amount of a nucleic acid encoding a p-Hyde protein, a nucleic acid encoding a fragment of p-Hyde protein, or the nucleic acid encoding a mutant p-Hyde protein and a pharmaceutical acceptable carrier or diluent, thereby suppressing cancer cells.

10 This invention provides a method of treating a subject with cancer which comprises administering to the subject a pharmaceutical composition comprising a therapeutically effective amount of a nucleic acid encoding a p-Hyde protein, a nucleic acid encoding a fragment of p-Hyde protein, or the nucleic acid encoding a mutant p-Hyde protein and a pharmaceutical acceptable carrier or diluent, thereby treating the subject with cancer.

15 This invention provides a method of treating a subject with cancer, comprising: 1) administering to the subject a pharmaceutical composition comprising a therapeutically effective amount of a nucleic acid encoding a p-Hyde protein, a nucleic acid encoding a fragment of p-Hyde protein, or the nucleic acid encoding a mutant p-Hyde protein in combination with radiation, chemotherapy, or UV mimetic drugs; and 2) a pharmaceutical  
20 acceptable carrier or diluent, thereby treating the subject with cancer.

The unique functional features of the *p-Hyde* gene may be exploited for the treatment of hyperproliferative disorders and cancer. One effective therapeutic strategy, for example, may be the treatment of carcinoma cells expressing *p-Hyde* with chemotherapy agents or UV  
25 mimetic drugs (such as acetylaminofluorine). However, cancer cells are not likely to produce significant levels of the growth inhibition *p-Hyde*. Consequently, the *p-Hyde* gene may be introduced into cancer cells by gene therapy. Tumors transduced with vectors containing *p-Hyde* may not only be directly suppressed by *p-Hyde* as demonstrated in this study, but also when treated in combination with DNA damaging therapy such as chemotherapy, UV  
30 mimetic drugs, or radiation, have even a greater anti-cancer effect. Since gene therapy will target cancer cells, then enhancement of apoptosis will occur more selectively in cancer cells following DNA damage (UV, radiation, or chemotherapy).

35 Three DNA enzyme repair systems were evaluated in parental compared to *p-Hyde* transfected cells: uridine phosphorylase, uridine kinase, and UV damage repair. UV damage



5 repair was impaired in the *p-Hyde* transfected cells. Figure 16 shows that decreased DNA  
repair activity results in higher levels of intact photoproducts (64PP). Consistent with these  
data, *pHyde* transfected cells also had a significant reduction in survival following UV  
exposure compared to parental AT3 cells as determined by colony formation assay. Thus,  
DNA repair enzyme impairment correlated with shorter survival and induction of apoptosis  
10 in prostate cancer cells transfected with *p-Hyde*.

This invention a method of treating a subject with cancer, comprising administering to the  
subject a therapeutically effective amount of a pharmaceutical composition comprising: 1)  
an adenovirus type 5 expression vector which comprises an adenovirus genome having a  
15 deletion in the E1 and E3 regions of the genome and an insertion within the regions of a full  
length sense p-Hyde cDNA under the control of a Rous Sarcoma virus promoter, and 2) a  
suitable carrier or diluent, thereby treating the subject with cancer. In one embodiment the  
cancer is selected from a group consisting of: melanoma; lymphoma; leukemia; and prostate,  
colorectal, pancreatic, breast, brain, or gastric carcinoma.

20 The present invention provides the means necessary for production of gene-based  
therapies directed at cancer cells. These therapeutic agents may take the form of  
polynucleotides comprising all or a portion of the p-Hyde locus placed in appropriate  
vectors or delivered to target cells in more direct ways such that the function of the p-  
Hyde protein is reconstituted. Therapeutic agents may also take the form of polypeptides  
25 based on either a portion of, or the entire protein sequence of p-Hyde. These may  
functionally replace the activity of p-Hyde in vivo.

A suitable bodily fluid includes, but is not limited to: serum, plasma, cerebrospinal fluid,  
30 lymphocytes, urine, transudates, or exudates. In the preferred embodiment, the suitable  
bodily fluid sample is serum or plasma. In addition, the bodily fluid sample may be cells  
from bone marrow, or a supernatant from a cell culture. Methods of obtaining a suitable  
bodily fluid sample from a subject are known to those skilled in the art. Methods of  
determining the level of antibody or antigen include, but are not limited to: ELISA, IFA, and  
35 Western blotting.

5 The diagnostic assays of the invention can be nucleic acid assays such as nucleic acid hybridization assays and assays which detect amplification of specific nucleic acid to detect for a nucleic acid sequence of the human p-Hyde described herein.

10 Target specific probes may be used in the nucleic acid hybridization diagnostic. The probes are specific for or complementary to the target of interest. For precise allelic differentiations, the probes should be about 14 nucleotides long and preferably about 20-30 nucleotides. For more general detection of the human p-Hyde of the invention, nucleic acid probes are about 50 to about 1000 nucleotides, most preferably about 200 to about 400 nucleotides.

15 The specific nucleic acid probe can be RNA or DNA polynucleotide or oligonucleotide, or their analogs. The probes may be single or double stranded nucleotides. The probes of the invention may be synthesized enzymatically, using methods well known in the art (*e.g.*, nick translation, primer extension, reverse transcription, the polymerase chain reaction, and others) or chemically (*e.g.*, by methods such as the phosphoramidite method described by  
20 Beaucage and Carruthers [19], or by the triester method according to Matteucci, *et al.* [62], both incorporated herein by reference).

5 An alternative means for determining the presence of the human p-Hyde is *in situ*  
hybridization, or more recently, *in situ* polymerase chain reaction. *In situ* PCR is described  
in Neuvo *et al.* [71], Intracellular localization of polymerase chain reaction (PCR)-amplified  
Hepatitis C cDNA; Bagasra *et al.* [10], Detection of Human Immunodeficiency virus type  
1 provirus in mononuclear cells by *in situ* polymerase chain reaction; and Heniford *et al.*  
10 [35], Variation in cellular EGF receptor mRNA expression demonstrated by *in situ* reverse  
transcriptase polymerase chain reaction. *In situ* hybridization assays are well known and are  
generally described in *Methods Enzymol.* [67] incorporated by reference herein. In an *in situ*  
hybridization, cells are fixed to a solid support, typically a glass slide. The cells are then  
contacted with a hybridization solution at a moderate temperature to permit annealing of  
15 target-specific probes that are labelled. The probes are preferably labelled with radioisotopes  
or fluorescent reporters.

The above described probes are also useful for in-situ hybridization or in order to locate  
tissues which express this gene, or for other hybridization assays for the presence of this gene  
20 or its MRNA in various biological tissues. In-situ hybridization is a sensitive localization  
method which is not dependent on expression of antigens or native vs. denatured conditions.

In brief, inhibitory nucleic acid therapy approaches can be classified into those that target  
25 DNA sequences, those that target RNA sequences (including pre-mRNA and mRNA), those  
that target proteins (sense strand approaches), and those that cause cleavage or chemical  
modification of the target nucleic acids.

Approaches targeting DNA fall into several categories. Nucleic acids can be designed to  
30 bind to the major groove of the duplex DNA to form a triple helical or "triplex" structure.  
Alternatively, inhibitory nucleic acids are designed to bind to regions of single stranded DNA  
resulting from the opening of the duplex DNA during replication or transcription.

More commonly, inhibitory nucleic acids are designed to bind to mRNA or mRNA  
35 precursors. Inhibitory nucleic acids are used to prevent maturation of pre-mRNA. Inhibitory

5 nucleic acids may be designed to interfere with RNA processing, splicing or translation.

The inhibitory nucleic acids can be targeted to mRNA. In this approach, the inhibitory nucleic acids are designed to specifically block translation of the encoded protein. Using this approach, the inhibitory nucleic acid can be used to selectively suppress certain cellular  
10 functions by inhibition of translation of mRNA encoding critical proteins. For example, an inhibitory nucleic acid complementary to regions of c-myc mRNA inhibits c-myc protein expression in a human promyelocytic leukemia cell line, HL60, which overexpresses the c-myc proto-oncogene. See Wickstrom E.L., *et al.* [93] and Harel-Bellan, A., *et al.* [31A]. As described in Helene and Toulme, inhibitory nucleic acids targeting mRNA have been  
15 shown to work by several different mechanisms to inhibit translation of the encoded protein(s).

Lastly, the inhibitory nucleic acids can be used to induce chemical inactivation or cleavage of the target genes or mRNA. Chemical inactivation can occur by the induction of crosslinks  
20 between the inhibitory nucleic acid and the target nucleic acid within the cell. Other chemical modifications of the target nucleic acids induced by appropriately derivatized inhibitory nucleic acids may also be used.

Cleavage, and therefore inactivation, of the target nucleic acids may be effected by attaching  
25 a substituent to the inhibitory nucleic acid which can be activated to induce cleavage reactions. The substituent can be one that affects either chemical, or enzymatic cleavage. Alternatively, cleavage can be induced by the use of ribozymes or catalytic RNA. In this approach, the inhibitory nucleic acids would comprise either naturally occurring RNA (ribozymes) or synthetic nucleic acids with catalytic activity.

30 As used herein, "pharmaceutical composition" could mean therapeutically effective amounts of polypeptide products of the invention together with suitable diluents, preservatives, solubilizers, emulsifiers, adjuvant and/or carriers useful in SCF (stem cell factor) therapy. A "therapeutically effective amount" as used herein refers to that amount which provides a  
35 therapeutic effect for a given condition and administration regimen. Such compositions are

5 liquids or lyophilized or otherwise dried formulations and include diluents of various buffer  
content (e.g., Tris-HCl, acetate, phosphate), pH and ionic strength, additives such as albumin  
or gelatin to prevent absorption to surfaces, detergents (e.g., Tween 20, Tween 80, Pluronic  
F68, bile acid salts), solubilizing agents (e.g., glycerol, polyethylene glycerol), anti-oxidants  
10 (e.g., ascorbic acid, sodium metabisulfite), preservatives (e.g., Thimerosal, benzyl alcohol,  
parabens), bulking substances or tonicity modifiers (e.g., lactose, mannitol), covalent  
attachment of polymers such as polyethylene glycol to the protein, complexation with metal  
ions, or incorporation of the material into or onto particulate preparations of polymeric  
compounds such as polylactic acid, polglycolic acid, hydrogels, etc, or onto liposomes,  
microemulsions, micelles, unilamellar or multilamellar vesicles, erythrocyte ghosts, or  
15 spheroplasts. Such compositions will influence the physical state, solubility, stability, rate  
of in vivo release, and rate of in vivo clearance of SCF. The choice of compositions will  
depend on the physical and chemical properties of the protein having SCF activity. For  
example, a product derived from a membrane-bound form of SCF may require a formulation  
containing detergent. Controlled or sustained release compositions include formulation in  
20 lipophilic depots (e.g., fatty acids, waxes, oils). Also comprehended by the invention are  
particulate compositions coated with polymers (e.g., poloxamers or poloxamines) and SCF  
coupled to antibodies directed against tissue-specific receptors, ligands or antigens or  
coupled to ligands of tissue-specific receptors. Other embodiments of the compositions of  
the invention incorporate particulate forms protective coatings, protease inhibitors or  
25 permeation enhancers for various routes of administration, including parenteral, pulmonary,  
nasal and oral. In one embodiment the pharmaceutical composition is administered  
parenterally, paracancerally, transmucosally, transdermally, intramuscularly, intravenously,  
intradermally, subcutaneously, intraperitoneally, intraventricularly, intracranially.

30 Further, as used herein "pharmaceutically acceptable carrier" are well known to those skilled  
in the art and include, but are not limited to, 0.01-0.1M and preferably 0.05M phosphate  
buffer or 0.8% saline. Additionally, such pharmaceutically acceptable carriers may be  
aqueous or non-aqueous solutions, suspensions, and emulsions. Examples of non-aqueous  
solvents are propylene glycol, polyethylene glycol, vegetable oils such as olive oil, and  
35 injectable organic esters such as ethyl oleate. Aqueous carriers include water,



5 alcoholic/aqueous solutions, emulsions or suspensions, including saline and buffered media. Parenteral vehicles include sodium chloride solution, Ringer's dextrose, dextrose and sodium chloride, lactated Ringer's or fixed oils. Intravenous vehicles include fluid and nutrient replenishers, electrolyte replenishers such as those based on Ringer's dextrose, and the like. Preservatives and other additives may also be present, such as, for example, antimicrobials,  
10 antioxidants, collating agents, inert gases and the like.

The term "adjuvant" refers to a compound or mixture that enhances the immune response to an antigen. An adjuvant can serve as a tissue depot that slowly releases the antigen and also as a lymphoid system activator that non-specifically enhances the immune response (Hood  
15 et al., *Immunology, Second Ed.*, 1984, Benjamin/Cummings: Menlo Park, California, p. 384). Often, a primary challenge with an antigen alone, in the absence of an adjuvant, will fail to elicit a humoral or cellular immune response. Adjuvant include, but are not limited to, complete Freund's adjuvant, incomplete Freund's adjuvant, saponin, mineral gels such as aluminum hydroxide, surface active substances such as lysolecithin, pluronic polyols,  
20 polyanions, peptides, oil or hydrocarbon emulsions, keyhole limpet hemocyanins, dinitrophenol, and potentially useful human adjuvant such as BCG (*bacille Calmette-Guerin*) and *Corynebacterium parvum*. Preferably, the adjuvant is pharmaceutically acceptable.

Controlled or sustained release compositions include formulation in lipophilic depots (e.g.  
25 fatty acids, waxes, oils). Also comprehended by the invention are particulate compositions coated with polymers (e.g. poloxamers or poloxamines) and the compound coupled to antibodies directed against tissue-specific receptors, ligands or antigens or coupled to ligands of tissue-specific receptors. Other embodiments of the compositions of the invention incorporate particulate forms protective coatings, protease inhibitors or permeation enhancers  
30 for various routes of administration, including parenteral, pulmonary, nasal and oral.

When administered, compounds are often cleared rapidly from mucosal surfaces or the circulation and may therefore elicit relatively short-lived pharmacological activity. Consequently, frequent administrations of relatively large doses of bioactive compounds may  
35 by required to sustain therapeutic efficacy. Compounds modified by the covalent attachment of water-soluble polymers such as polyethylene glycol, copolymers of polyethylene glycol

5 and polypropylene glycol, carboxymethyl cellulose, dextran, polyvinyl alcohol, polyvinylpyrrolidone or polyproline are known to exhibit substantially longer half-lives in blood following intravenous injection than do the corresponding unmodified compounds (Abuchowski et al., 1981; Newmark et al., 1982; and Katre et al., 1987). Such modifications may also increase the compound's solubility in aqueous solution, eliminate aggregation, 10 enhance the physical and chemical stability of the compound, and greatly reduce the immunogenicity and reactivity of the compound. As a result, the desired *in vivo* biological activity may be achieved by the administration of such polymer-compound adducts less frequently or in lower doses than with the unmodified compound.

15 *Dosages.* The sufficient amount may include but is not limited to from about 1  $\mu\text{g/kg}$  to about 1000 mg/kg. The amount may be 10 mg/kg. The pharmaceutically acceptable form of the composition includes a pharmaceutically acceptable carrier.

The preparation of therapeutic compositions which contain an active component is well 20 understood in the art. Typically, such compositions are prepared as an aerosol of the polypeptide delivered to the nasopharynx or as injectables, either as liquid solutions or suspensions, however, solid forms suitable for solution in, or suspension in, liquid prior to injection can also be prepared. The preparation can also be emulsified. The active therapeutic ingredient is often mixed with excipients which are pharmaceutically acceptable 25 and compatible with the active ingredient. Suitable excipients are, for example, water, saline, dextrose, glycerol, ethanol, or the like and combinations thereof. In addition, if desired, the composition can contain minor amounts of auxiliary substances such as wetting or emulsifying agents, pH buffering agents which enhance the effectiveness of the active ingredient.

30 An active component can be formulated into the therapeutic composition as neutralized pharmaceutically acceptable salt forms. Pharmaceutically acceptable salts include the acid addition salts (formed with the free amino groups of the polypeptide or antibody molecule) and which are formed with inorganic acids such as, for example, hydrochloric or phosphoric 35 acids, or such organic acids as acetic, oxalic, tartaric, mandelic, and the like. Salts formed

5 from the free carboxyl groups can also be derived from inorganic bases such as, for example, sodium, potassium, ammonium, calcium, or ferric hydroxides, and such organic bases as isopropylamine, trimethylamine, 2-ethylamino ethanol, histidine, procaine, and the like.

A composition comprising "A" (where "A" is a single protein, DNA molecule, vector, etc.)  
10 is substantially free of "B" (where "B" comprises one or more contaminating proteins, DNA molecules, vectors, etc.) when at least about 75% by weight of the proteins, DNA, vectors (depending on the category of species to which A and B belong) in the composition is "A". Preferably, "A" comprises at least about 90% by weight of the A+B species in the composition, most preferably at least about 99% by weight.

15 The phrase "therapeutically effective amount" is used herein to mean an amount sufficient to reduce by at least about 15 percent, preferably by at least 50 percent, more preferably by at least 90 percent, and most preferably prevent, a clinically significant deficit in the activity, function and response of the host.

20 According to the invention, the component or components of a therapeutic composition of the invention may be introduced parenterally, transmucosally, *e.g.*, orally, nasally, pulmonarailly, or rectally, or transdermally. Preferably, administration is parenteral, *e.g.*, via intravenous injection, and also including, but is not limited to, intra-arteriole, intramuscular,  
25 intradermal, subcutaneous, intraperitoneal, intraventricular, and intracranial administration. Oral or pulmonary delivery may be preferred to activate mucosal immunity; since pneumococci generally colonize the nasopharyngeal and pulmonary mucosa, mucosal immunity may be a particularly effective preventive treatment. The term "unit dose" when used in reference to a therapeutic composition of the present invention refers to physically  
30 discrete units suitable as unitary dosage for humans, each unit containing a predetermined quantity of active material calculated to produce the desired therapeutic effect in association with the required diluent; *i.e.*, carrier, or vehicle.

According to the present invention, a method is also provided of supplying wild-type  
35 p-Hyde function to a cell which carries mutant p-Hyde alleles. Supplying such a function

5 should suppress neoplastic growth of the recipient cells. The wild-type p-Hyde gene or a  
part of the gene may be introduced into the cell in a vector such that the gene remains  
extrachromosomal. In such a situation, the gene will be expressed by the cell from the  
extrachromosomal location. If a gene fragment is introduced and expressed in a cell  
10 carrying a mutant p-Hyde allele, the gene fragment should encode a part of the p-Hyde  
protein which is required for non-neoplastic growth of the cell. More preferred is the  
situation where the wild-type p-Hyde gene or a part thereof is introduced into the mutant  
cell in such a way that it recombines with the endogenous mutant p-Hyde gene present in  
the cell. Such recombination requires a double recombination event which results in the  
correction of the p-Hyde gene mutation. Vectors for introduction of genes both for  
15 recombination and for extrachromosomal maintenance are known in the art, and any  
suitable vector may be used. Methods for introducing DNA into cells such as  
electroporation, calciumphosphate co-precipitation and viral transduction are known in the  
art, and the choice of method is within the competence of the routineer. Cells transformed  
with the wild-type p-Hyde gene can be used as model systems to study cancer remission and  
20 drug treatments which promote such remission.

As generally discussed above, the p-Hyde gene or fragment, where applicable, may  
be employed in gene therapy methods in order to increase the amount of the expression  
products of such genes in cancer cells. Such gene therapy is particularly appropriate for  
25 use in both cancerous and pre-cancerous cells, in which the level of p-Hyde polypeptide is  
absent or diminished compared to normal cells. It may also be useful to increase the level  
of expression of a given p-Hyde gene even in those tumor cells in which the mutant gene  
is expressed at a "normal" level, but the gene product is not fully functional.

30 Gene therapy would be carried out according to generally accepted methods, for  
example, as described by Friedman, 1991. Cells from a patient's tumor would be first  
analyzed by the diagnostic methods described above, to ascertain the production of p-Hyde  
polypeptide in the tumor cells. A virus or plasmid vector (see further details below),  
containing a copy of the p-Hyde gene linked to expression control elements and capable  
35 of replicating inside the tumor cells, is prepared. Suitable vectors are known, such as

5 disclosed in U.S. Pat. No. 5,252,479 and PCT published application WO 93/07282. The vector is then injected into the patient, either locally at the site of the tumor or systemically (in order to reach any tumor cells that may have metastasized to other sites). If the transfected gene is not permanently incorporated into the genome of each of the targeted minor cells, the treatment may have to be repeated periodically.

10 Gene transfer systems known in the art may be useful in the practice of the gene therapy methods of the present invention. These include viral and nonviral transfer methods. A number of viruses have been used as gene transfer vectors, including papovaviruses, e.g., SV40 (Madzak et al., 1992), adenovirus (Berkner, 1992; Berkner et al., 1988; Gorziglia and Kapikian, 1992; Quantin et al., 1992; Rosenfeld et al., 1992; Wilkinson et al., 1992; 15 Stratford-Perricaudet et al., 1990), vaccinia virus (Moss, 1992), adeno-associated virus (Muzyczka, 1992; Ohi et al., 1990), herpesviruses including HSV and EBV (Margolskee, 1992; Johnson et al., 1992; Fink et al., 1992; Breakfield and Geller, 1987; Freese et al., 1990), and retroviruses of avian (Bradyopadhyay and Temin, 1984; Petropoulos et al., 1992), murine (Miller, 1992; Miller et al., 1985; Sorge et al., 1984; Mann and Baltimore, 1985; Miller et al., 1988), and human origin (Shimada et al., 1991; Helseth et al., 1990; Page et al., 1990; Buchschacher and Panganiban, 1992). Most human gene 20 therapy protocols have been based on disabled murine retroviruses.

Peptides which have p-Hyde activity can be supplied to cells which carry mutant or 25 missing p-Hyde alleles. The sequence of the p-Hyde protein is disclosed (SEQ ID NO:2). Protein can be produced by expression of the cDNA sequence in bacteria, for example, using known expression vectors. Alternatively, p-Hyde polypeptide can be extracted from p-Hyde-producing mammalian cells. In addition, the techniques of synthetic chemistry can be employed to synthesize p-Hyde protein. Any of such techniques can provide the 30 preparation of the present invention which comprises the p-Hyde protein. The preparation is substantially free of other human proteins. This is most readily accomplished by synthesis in a microorganism or in vitro.

P-Hyde molecules can be introduced into cells by microinjection or by use of liposomes, for 35 example. Alternatively, some active molecules may be taken up by cells, actively or by



5 diffusion. Extracellular application of the p-Hyde gene product may be sufficient to affect  
tumor growth. Supply of molecules with p-Hyde activity should lead to partial reversal of  
the neoplastic state. Other molecules with p-Hyde activity (for example, peptides, drugs  
or organic compounds) may also be used to effect such a reversal. Modified polypeptides  
having substantially similar function are also used for peptide therapy.

10

In another embodiment, the active compound can be delivered in a vesicle, in particular a  
liposome (see Langer, Science 249:1527-1533 (1990); Treat et al., in *Liposomes in the  
Therapy of Infectious Disease and Cancer*, Lopez-Berestein and Fidler (eds.), Liss, New  
York, pp. 353-365 (1989); Lopez-Berestein, *ibid.*, pp. 317-327; see generally *ibid.*).

15

In yet another embodiment, the therapeutic compound can be delivered in a controlled  
release system. For example, the polypeptide may be administered using intravenous  
infusion, an implantable osmotic pump, a transdermal patch, liposomes, or other modes of  
administration. In one embodiment, a pump may be used (see Langer, *supra*; Sefton, CRC  
20 Crit. Ref. Biomed. Eng. 14:201 (1987); Buchwald et al., Surgery 88:507 (1980); Saudek et  
al., N. Engl. J. Med. 321:574 (1989)). In another embodiment, polymeric materials can be  
used (see *Medical Applications of Controlled Release*, Langer and Wise (eds.), CRC Pres.,  
Boca Raton, Florida (1974); *Controlled Drug Bioavailability, Drug Product Design and  
Performance*, Smolen and Ball (eds.), Wiley, New York (1984); Ranger and Peppas, J.  
25 Macromol. Sci. Rev. Macromol. Chem. 23:61 (1983); see also Levy et al., Science 228:190  
(1985); During et al., Ann. Neurol. 25:351 (1989); Howard et al., J. Neurosurg. 71:105  
(1989)). In yet another embodiment, a controlled release system can be placed in proximity  
of the therapeutic target, i.e., the brain, thus requiring only a fraction of the systemic dose  
(see, e.g., Goodson, in *Medical Applications of Controlled Release*, *supra*, vol. 2, pp. 115-  
30 138 (1984)). Preferably, a controlled release device is introduced into a subject in proximity  
of the site of inappropriate immune activation or a tumor. Other controlled release systems  
are discussed in the review by Langer (Science 249:1527-1533 (1990)).

35

A subject in whom administration of an active component as set forth above is an effective  
therapeutic regimen for a bacterial infection is preferably a human, but can be any animal.

5 Thus, as can be readily appreciated by one of ordinary skill in the art, the methods and  
pharmaceutical compositions of the present invention are particularly suited to administration  
to any animal, particularly a mammal, and including, but by no means limited to, domestic  
animals, such as feline or canine subjects, farm animals, such as but not limited to bovine,  
equine, caprine, ovine, and porcine subjects, wild animals (whether in the wild or in a  
10 zoological garden), research animals, such as mice, rats, rabbits, goats, sheep, pigs, dogs,  
cats, etc., *i.e.*, for veterinary medical use.

15 In the therapeutic methods and compositions of the invention, a therapeutically effective  
dosage of the active component is provided. A therapeutically effective dosage can be  
determined by the ordinary skilled medical worker based on patient characteristics (age,  
weight, sex, condition, complications, other diseases, etc.), as is well known in the art.  
Furthermore, as further routine studies are conducted, more specific information will emerge  
regarding appropriate dosage levels for treatment of various conditions in various patients,  
and the ordinary skilled worker, considering the therapeutic context, age and general health  
20 of the recipient, is able to ascertain proper dosing. Generally, for intravenous injection or  
infusion, dosage may be lower than for intraperitoneal, intramuscular, or other route of  
administration. The dosing schedule may vary, depending on the circulation half-life, and  
the formulation used. The compositions are administered in a manner compatible with the  
dosage formulation in the therapeutically effective amount. Precise amounts of active  
25 ingredient required to be administered depend on the judgment of the practitioner and are  
peculiar to each individual. However, suitable dosages may range from about 0.1 to 20,  
preferably about 0.5 to about 10, and more preferably one to several, milligrams of active  
ingredient per kilogram body weight of individual per day and depend on the route of  
administration. Suitable regimes for initial administration and booster shots are also  
30 variable, but are typified by an initial administration followed by repeated doses at one or  
more hour intervals by a subsequent injection or other administration. Alternatively,  
continuous intravenous infusion sufficient to maintain concentrations of ten nanomolar to  
ten micromolar in the blood are contemplated.

35 It is contemplated by this invention that p-Hyde replacement therapy could be used similarly.

5 in conjunction with chemo- or radiotherapeutic intervention. To induce susceptibility to  
cell death or to inhibit cell growth or to kill cells, such as malignant or metastatic cells, using  
the methods and compositions of the present invention, one would contact a "target" cell with  
the expression vector and at least one DNA damaging agent. In one embodiment the cell is  
10 contacted with a single composition or pharmacological formulation that includes both  
agents, or by contacting the cell with two distinct compositions or formulations, at the same  
time, wherein one composition includes the vector and the other includes the DNA damaging  
agent. In another embodiment, treatment with the vector may precede or follow the DNA  
damaging agent treatment by intervals ranging from minutes to weeks. Protocols and  
methods are known to those skilled in the art.

15 DNA damaging agents or factors are known to those skilled in the art and means any  
chemical compound or treatment method that induces DNA damage when applied to a cell.  
Such agents and factors include radiation and waves that induce DNA damage such as,  
gamma -irradiation, X-rays, UV-irradiation, microwaves, electronic emissions, and the like.  
20 A variety of chemical compounds, also described as "chemotherapeutic agents", function to  
induce DNA damage, all of which are intended to be of use in the combined treatment  
methods disclosed herein. Chemotherapeutic agents contemplated to be of use, include, e.g.,  
adriamycin, 5-fluorouracil (5FU), etoposide (VP-16), camptothecin, actinomycin-D,  
mitomycin C, cisplatin (CDDP) and even hydrogen peroxide. The invention also  
25 encompasses the use of a combination of one or more DNA damaging agents, whether  
radiation-based or actual compounds, such as the use of X-rays with cisplatin or the use of  
cisplatin with etoposide.

30 In another embodiment one may irradiate the localized tumor site with DNA damaging  
radiation such as X-rays, UV-light, gamma -rays or even microwaves. Alternatively, the  
tumor cells may be contacted with the DNA damaging agent by administering to the subject  
a therapeutically effective amount of a pharmaceutical composition comprising a DNA  
damaging compound such as, adriamycin, 5-fluorouracil, etoposide, camptothecin,  
actinomycin-D, mitomycin C, or more preferably, cisplatin. The DNA damaging agent may  
35 be prepared and used as a combined therapeutic composition, or kit, by combining it with a

5 p-Hyde expression construct, as described above. Agents that directly cross-link nucleic acids, specifically DNA, are envisaged and are shown herein, to eventuate DNA damage leading to a synergistic antineoplastic combination. Agents such as cisplatin, and other DNA alkylating may be used. Cisplatin has been widely used to treat cancer, with efficacious doses used in clinical applications of  $20 \text{ mg/m}^2$  for 5 days every three weeks for a total of three courses. Cisplatin is not absorbed orally and must therefore be delivered via injection intravenously, subcutaneously, intratumorally or intraperitoneally. Agents that damage DNA also include compounds that interfere with DNA replication, mitosis and chromosomal segregation. Such chemotherapeutic compounds include adriamycin, also known as doxorubicin, etoposide, verapamil, podophyllotoxin, and the like. Widely used in a clinical setting for the treatment of neoplasms, these compounds are administered through bolus injections intravenously at doses ranging from  $25\text{-}75 \text{ mg/m}^2$  at 21 day intervals for adriamycin, to  $35\text{-}50 \text{ mg/m}^2$  for etoposide intravenously or double the intravenous dose orally.

20 Agents that disrupt the synthesis and fidelity of nucleic acid precursors and subunits also lead to DNA damage. As such a number of nucleic acid precursors have been developed. Particularly useful are agents that have undergone extensive testing and are readily available. As such, agents such as 5-fluorouracil (5-FU), are preferentially used by neoplastic tissue, making this agent particularly useful for targeting to neoplastic cells. Although quite toxic, 5-FU, is applicable in a wide range of carriers, including topical, however intravenous administration with doses ranging from 3 to 15 mg/kg/day being commonly used.

30 Other factors that cause DNA damage and have been used extensively include what are commonly known as gamma -rays, X-rays, and/or the directed delivery of radioisotopes to tumor cells. Other forms of DNA damaging factors are also contemplated such as microwaves and UV-irradiation. It is most likely that all of these factors effect a broad range of damage DNA, on the precursors of DNA, the replication and repair of DNA, and the assembly and maintenance of chromosomes. Dosage ranges for X-rays range from daily doses of 50 to 200 roentgens for prolonged periods of time (3 to 4 weeks), to single doses of 2000 to 6000 roentgens. Dosage ranges for radioisotopes vary widely, and depend on the

5 half-life of the isotope, the strength and type of radiation emitted, and the uptake by the neoplastic cells.

Several non-viral methods for the transfer of expression constructs into cultured mammalian cells also are contemplated by the present invention. These include calcium phosphate  
10 precipitation (Graham and Van Der Eb, 1973; Chen and Okayama, 1987; Rippe et al., 1990) DEAE-dextran (Gopal, 1985), electroporation (Tur-Kaspa et al., 1986; Potter et al., 1984), direct microinjection (Harland and Weintraub, 1985), DNA-loaded liposomes (Nicolau and Sene, 1982; Fraley et al., 1979) and lipofectamine-DNA complexes, cell sonication  
15 (Fechheimer et al., 1987), gene bombardment using high velocity microprojectiles (Yang et al., 1990), and receptor-mediated transfection (Wu and Wu, 1987; Wu and Wu, 1988). Some of these techniques may be successfully adapted for in vivo or ex vivo use. Also, helper cell lines may be derived from human cells such as human embryonic kidney cells, muscle cells, hematopoietic cells or other human embryonic mesenchymal or epithelial cells. Alternatively, the helper cells may be derived from the cells of other mammalian species that  
20 are permissive for human adenovirus. Such cells include, e.g., Vero cells or other monkey embryonic mesenchymal or epithelial cells. As stated above, the preferred helper cell line is 293.

In another embodiment, the active compound can be delivered in a vesicle, in particular a  
25 liposome (see Langer, Science 249:1527-1533 (1990); Treat et al., in *Liposomes in the Therapy of Infectious Disease and Cancer*, Lopez-Berestein and Fidler (eds.), Liss, New York, pp. 353-365 (1989); Lopez-Berestein, *ibid.*, pp. 317-327; see generally *ibid.*).

In yet another embodiment, the therapeutic compound can be delivered in a controlled  
30 release system. For example, the polypeptide may be administered using intravenous infusion, an implantable osmotic pump, a transdermal patch, liposomes, or other modes of administration. In one embodiment, a pump may be used (see Langer, *supra*; Sefton, CRC Crit. Ref. Biomed. Eng. 14:201 (1987); Buchwald et al., Surgery 88:507 (1980); Saudek et al., N. Engl. J. Med. 321:574 (1989)). In another embodiment, polymeric materials can be  
35 used (see *Medical Applications of Controlled Release*, Langer and Wise (eds.), CRC Pres.,



5 Boca Raton, Florida (1974); *Controlled Drug Bioavailability, Drug Product Design and Performance*, Smolen and Ball (eds.), Wiley, New York (1984); Ranger and Peppas, J. Macromol. Sci. Rev. Macromol. Chem. 23:61 (1983); see also Levy et al., Science 228:190 (1985); During et al., Ann. Neurol. 25:351 (1989); Howard et al., J. Neurosurg. 71:105 (1989)). In yet another embodiment, a controlled release system can be placed in proximity  
10 of the therapeutic target, i.e., the brain, thus requiring only a fraction of the systemic dose (see, e.g., Goodson, in Medical Applications of Controlled Release, supra, vol. 2, pp. 115-138 (1984)). Preferably, a controlled release device is introduced into a subject in proximity of the site of inappropriate immune activation or a tumor. Other controlled release systems are discussed in the review by Langer (Science 249:1527-1533 (1990)).

15 As can be readily appreciated by one of ordinary skill in the art, the methods and pharmaceutical compositions of the present invention are particularly suited to administration to a mammal, preferable a human subject. In the therapeutic methods and compositions of the invention, a therapeutically effective dosage of the active component is provided. A  
20 therapeutically effective dosage can be determined by the ordinary skilled medical worker based on patient characteristics (age, weight, sex, condition, complications, other diseases, etc.), as is well known in the art. Furthermore, as further routine studies are conducted, more specific information will emerge regarding appropriate dosage levels for treatment of various conditions in various patients, and the ordinary skilled worker, considering the therapeutic  
25 context, age and general health of the recipient, is able to ascertain proper dosing. Generally, for intravenous injection or infusion, dosage may be lower than for intraperitoneal, intramuscular, or other route of administration. The dosing schedule may vary, depending on the circulation half-life, and the formulation used. The compositions are administered in a manner compatible with the dosage formulation in the therapeutically effective amount.  
30 Precise amounts of active ingredient required to be administered depend on the judgment of the practitioner and are peculiar to each individual. However, suitable dosages may range from about 0.1 to 20, preferably about 0.5 to about 10, and more preferably one to several, milligrams of active ingredient per kilogram body weight of individual per day and depend on the route of administration. Suitable regimes for initial administration and booster shots  
35 are also variable, but are typified by an initial administration followed by repeated doses at

5 one or more hour intervals by a subsequent injection or other administration. Alternatively, continuous intravenous infusion sufficient to maintain concentrations of ten nanomolar to ten micromolar in the blood are contemplated.

10 The present invention provides a kit comprising the all the essential materials and reagents required for inhibiting prostate tumor cell proliferation, transforming prostate cells or detecting prostate cancer cells, may be assembled together in a kit. This generally will comprise selected expression constructs. Also included may be various media for replication of the expression constructs and host cells for such replication. Such kits will comprise distinct containers for each individual reagent. When the components of the kit are provided 15 in one or more liquid solutions, the liquid solution preferably is an aqueous solution, with a sterile aqueous solution being particularly preferred. For in vivo use, the expression construct may be formulated into a pharmaceutically acceptable syringeable composition. In this case, the container means may itself be an inhalent, syringe, pipette, eye dropper, or other such like apparatus, from which the formulation may be applied to an infected area of the 20 body, such as the lungs, injected into an animal, or even applied to and mixed with the other components of the kit. The components of the kit may also be provided in dried or lyophilized forms. When reagents or components are provided as a dried form, reconstitution generally is by the addition of a suitable solvent. It is envisioned that the solvent also may be provided in another container means.

25 The kits of the present invention also will typically include a means for containing the vials in close confinement for commercial sale such as, e.g., injection or blow-molded plastic containers into which the desired vials are retained. Irrespective of the number or type of containers, the kits of the invention also may comprise, or be packaged with, an instrument 30 for assisting with the injection/administration or placement of the ultimate complex composition within the body of an animal. Such an instrument may be an inhalent, syringe, pipette, forceps, measured spoon, eye dropper or any such medically approved delivery vehicle.

35 The following examples are presented in order to more fully illustrate the preferred

5       embodiments of the invention. They should in no way be construed, however, as limiting  
the broad scope of the invention.

## EXPERIMENTAL DETAILS SECTION

10       **EXAMPLE: p-Hyde induces susceptibility to induction of cell programmed cell death  
in prostate cancer**

### MATERIALS AND METHODS

15       **Cell lines:** Two rat prostatic cancer cell lines (anaplastic tumor, low metastatic AT-1 and  
highly metastatic MAT-LyLu) were used for the cDNA competition hybridization strategy  
(Rinaldy and Steiner, 1997). Other Dunning rat prostate cancer cell lines used for Northern  
analyses were AT-3, MAT-LyLu, MAT-Lu and G. These cell lines were derived from  
established *in-vivo* Dunning R3327 rat prostatic tumor sublines and further developed as *in-*  
*vitro* cell lines by Isaacs *et al.* at Johns Hopkins Oncology Center, Baltimore, Maryland  
20       (Isaacs et al., 1986). Human prostate cancer cell lines PPC-1, LNCaP, TSU and DU145 used  
for Northern analyses were obtained from American Tissue Culture Collection, Rockville,  
Maryland. All cells were grown in RPMI 1640 medium (Mediatech, Herndon, Virginia) in  
the presence of 10% fetal calf serum, 50 units of penicillin G and 50 µg streptomycin sulfate  
per ml and 250 µM dexamethasone as previously described (Isaacs et al, 1986).

25       **Cloning strategy:** Radiolabeled MAT-LyLu cDNA population in the presence of vast excess  
amount of competitor non-radiolabeled AT-1 cDNA population was used to identify cDNAs  
clones in the MAT-LyLu cDNA library (Rinaldy and Steiner, 1997). One of these cDNAs  
was novel and designated as *p-Hyde*. The prostate cancer associated *p-Hyde* cDNA was  
30       further characterized.

35       **Characterization of cDNA: Sequencing-** *p-Hyde* cDNA was originally obtained as a ? ZAP  
Uni XR clone, and was further subcloned into pBluescript SK<sup>-</sup> vector through *in vivo*  
excision protocol as described (Stratagene, La Jolla, California). This double-stranded  
cDNA was further subjected for Dye Terminator Cycle Sequencing (Perkin Elmer, Foster

5 City, California) using ABI 377 automatic DNA sequencer Version 3.0. The open reading  
frame of *p-Hyde* cDNA was determined using the DNA Strider program (Pasteur Institute,  
Paris).

10 **Northern blot analyses:** Total RNAs were isolated from cell lines during exponential  
growth (70% confluence) by using RNAzol B as suggested by the supplier (Tel-Test, Inc.,  
Friendswood, Texas) and subjected to Northern blot analyses. Reverse and forward PCR  
primer were designed and used to amplify 1.35 kb of the *p-Hyde* sequence representing the  
1467 bases of its open reading frame. The PCR product was radiolabeled by random oligo  
15 labeling technique (Multiprime DNA Labeling System, Amersham) and used as a probe for  
Northern analyses. Relative expression of *p-Hyde* gene was compared to an internal control  
(cyclophylin) by scanning of the autoradiogram with a PDI Discovery Series Scanner  
equipped with Quantity One software.

20 **In vitro transcription and translation:** The open reading frame of the cDNA was confirmed  
using *in-vitro* transcription (capped mRNA) as described followed by *in-vitro* translation  
using rabbit reticulocyte lysate (Stratagene, La Jolla, California) in the presence of 75 Ci of  
L-[<sup>35</sup>S] Methionine (Amersham, Arlington Heights, Illinois). The *in vitro* translation  
product was further identified on 10% SDS polyacrylamide gel electrophoresis by  
autoradiography.

25 **Subcloning of *p-Hyde* into pcDNA3.1 (-):** cDNA insert was released from the pBluescript  
SK<sup>-</sup> vector through double digests by KpnI and SacI (SK fragment). This fragment  
represents the intact *p-Hyde* sequence and was then ligated into dephosphorylated KpnI-SacI  
double digests of mammalian shuttle vector pcDNA 3.1 (-) (Invitrogen). Ligation mix was  
30 used to transfect competent DH5<sup>+</sup>, selected for ampicillin resistance followed by plasmid  
preparation using standard cesium chloride density gradient centrifugation. The new  
construct of *p-Hyde* was then used to transfect AT3 rat prostatic cancer subline by using  
lipofectamine (Gibco/BRL) followed by G418 selection (Rinaldy et al., 1988). Eight clones  
were obtained and two of them, AT3-H1 and AT3-H2, were used to assess the function of  
35 *p-Hyde* in its association with apoptosis. In addition, AT3 cell line was also transfected with

5 pcDNA3.1(-) vector only and its stable transfected cell line, designated as AT3-pc was used as negative control relative to stable transfectant of AT3-H1 and AT3-H2 for the functional assessment of the *p-Hyde*.

10 **Apoptosis assay:** Apoptosis intensity in AT3 parental cell line was assessed in comparison with AT3-H1, AT3-H2 and AT3-pc as negative control. Two apoptotic agents were employed for this assessment: (1) UV damage of the DNA using UV dosage of 200 J/m<sup>2</sup>; apoptosis intensity was assayed at 36 h post-UV irradiation and (2) 100  $\mu$ M Fluorodeoxyuridine (FUrD) treatment for 36 hours followed by the apoptosis assay. After these apoptotic induction, cells were collected in two fractions: floating and attached cells.  
15 Both cell fractions were counted using Neubauer chamber and trypan blue exclusion. DNA were extracted separately from both fractions and analyzed on 1.6 % agarose gel electrophoresis to visualize the DNA laddering.

**UV-damage repair assay:** Ultraviolet induced damage in the DNA was assayed using mouse monoclonal antibody specifically cross reacts with cyclobutane pyrimidine dimer (TDM-2) and photoproduct (64M-2). The presence of cyclobutane dimer and 64 photoproducts in DNAs were assessed in microtiter plates (100 and 200 ng/well) using both antibodies separately in a standard ELISA technique. In addition, the UV resistance was also assessed by using UV gradient assay as published (Rinaldy et al, 1988).  
20

25 **Uridine Phosphorylase Assay:** Cell extract will be prepared from the cell pellet Before and after the induction with 1 mM 5-dFUrD for 24 hours, the corresponding cell extract will be prepared in 50 mM Potassium Phosphate buffer pH 7.4 through sonication followed by dialysis against the reaction buffer (50 mM potassium phosphate, pH 7.4). The amount of protein will be determined by using standard Lowry or Biorad assay. The same amount of protein from all cell lines will be assayed for UP activity in 50 mM potassium phosphate  
30 buffer (pH 7.4) containing 10 mM uridine or thymidine as substrate. After 30 min incubation at 37° C, the reaction will be terminated by adding methanol followed by centrifugation. An aliquot of the supernatant will be run on HPLC column (6 x 200 mm) of ERC-ODS-1171 (ERMA CR, Inc). The amount of reaction product Uracil or Thymidine can be measured



5 with UV detector at 265 nm compared to the standard. As negative control, the similar reaction mixture will be boiled before the incubation.

10 **Construction of cDNA libraries:** In the first stage, cDNA libraries derived from AT-1 and MAT-LyLu cell lines were generated using Uni ZAP XR vector based on the protocol of Stratagene. The independent clones obtained were 1.9 and 3.4 million clones for MAT-LyLu and AT-1, respectively. These unamplified libraries were subjected to PCR amplification of the cDNA insert population. Reverse primers (RP) and forward primers (FP), downstream and upstream of XhoI and Eco RI cloning site, were used to amplify the cDNA insert population. The distance between both primers in ? Uni ZAP or pBluescript was 228 bases.

15 **Design of competition probes.** Two PCR probes were amplified: radiolabeled MAT-LyLu cDNA population probe and the non-radiolabeled AT-1 cDNA population probe (the cold competitor). The radiolabeled MAT-LyLu PCR product was enriched using S400 Sephacryl spin column. The majority of the unincorporated <sup>32</sup>P-dCTP, primer dimer, and 228 bp of PCR product resulting from the amplification of ? DNA without insert, was separated from the cDNAs. The purified radiolabeled cDNAs were mixed with 30 fold excess of non-radiolabeled cold competitor AT-1 PCR products and used as a competition-probe to screen the MAT-LyLu cDNA library.

25 Two kinds of unexpected radiolabeled PCR products that may potentially interfere with the hybridization between the radiolabeled cDNA of the competition probe and the screened cDNA of the library were: 1) the non-exponential amplification of the cDNAs, and 2) 228 bp PCR product derived from the ? DNA without cDNA insert. In order to reduce the possible cross-hybridization between these two unexpected PCR products with the vector of the screened library, excess amounts of HindIII-digested \_DNA, PvuII-digested-pBluescript DNA, and the 228 bp PCR product of the pBluescript based on both primers were mixed with the competition probe. Preliminary assessment of this complete mixed competition probe indicated that the hybridization of the MAT-LyLu cDNA library with this probe was extremely weak; whereas the duplicate filter hybridized with the same probe, but without non-radiolabeled AT-1 PCR product, was extremely positive. This clearly indicate that the

5 positive hybridization of the MAT-LyLu products was due to the radiolabeled MAT-LyLu  
cDNAs of the PCR products which was not competed by the AT-1 cDNAs.

10 **Screening of MAT-LyLu cDNA Libraries.** Independent cDNA clones of the unamplified  
MAT-LyLu phage cDNA library (250,000 clones) were screened with the competition probe  
as described above. Nineteen enhanced signals were observed; these putative prostate cancer-  
or metastasis-associated plaques were then rescreened. As the result of this rescreening, 12  
individual plaques were purified. The cDNAs were excised and subcloned into plasmid  
based vectors (pBluescript). This is possible due to the ability of helper phage ExAssist  
(Stratagene) to excise the cDNA with the pBluescript sequence in circular form as a  
15 filamentous phage and secreted from the cell (Fig. 2). Recombinant pBluescript plasmids  
were recovered by infecting an F' strain of E. coli (SOLR strain, Stratagene), ampicillin  
resistant colonies were selected, and the plasmid DNA was extracted. The obtained plasmids  
were digested with Eco RI and XhoI to identify the cDNA inserts by agarose gel  
electrophoresis. The results indicated that only four plasmids carried cDNA inserts.

20 **Sequencing of four candidate genes:** All four cDNAs were partially sequenced at both the  
3' and 5'-ends. These initial sequences were used to search the Genbank nucleic acid  
database of NCBI for homology or similarities. Three of the cDNAs matched known  
sequences: 1) rat mitochondrial genes coding for 16s and 12s rRNAs and tRNAs specific for  
25 valine and phenylalanine (Accession # emb/V00680/MIRNR2). 2) rat nucleolar proteins  
B23.1 mRNA (Accession # gb/J03969/RATB23NP) and 3) rat nucleolar proteins B23.1 and  
B23.2 (Accession # gb/M37041/ RATNUCBA7). The results of this competitive  
hybridization were consistent with the recognized phenotypic difference between both cell  
lines. MAT-LyLu exhibits twice the number of both nucleoli and mitochondria (Isaacs et al,  
30 1986). This also suggests that the MAT-LyLu cell line is metabolically more active due to  
higher gene-dosage or gene-amplification of nucleolar and mitochondrial rRNA and its  
associated genes.

35 **The complete sequence of the putative gene, *p-Hyde*:** The fourth cDNA. (lane 1),  
designated as *p-Hyde*, had an initial sequence which did not match any known full length

5 sequences in the BLASTN nucleic acid database of the NCBI. Accordingly, *p-Hyde* cDNA  
was completely sequenced by using a walk-through sequencing strategy in both directions  
for three redundancies. Nine contigs from each sequencing direction were obtained and  
compounded as a full length composite of 2694 bases. The poly(A) tail on the 3'-end and  
polyadenylation signal sequence of GAGAAA (a slight modification of the conserved  
10 AATAAA sequence) located at the position 27 upstream from the poly(A) was also  
identified. Fig.6 shows the sequencing strategy and the restriction map illustrating the open  
reading frame of *p-Hyde*. SEQ ID No. 3 sets forth the resulting nucleic acid sequence of the  
rat *p-Hyde* and the resulting amino acid sequence of the rat *p-Hyde*.

15 **Generation of PCR probe representing open reading frame *p-Hyde* Gene:** Figure 8  
shows the PCR amplification of the cDNA insert using three sets of sequencing primers.  
PCR product based on primer 11 and 5 represents the open reading frame of the *p-Hyde*  
cDNA . It was further radiolabeled by Random-Oligo-Labeling technique and used as a  
representative probe for the cDNA insert in Northern analysis.

20 **Northern blot analysis of *p-Hyde*.** Some of the rat prostate cancer sublines (AT-1, MAT-  
LyLu, MAT-Lu, AT-3 and G) and human prostate cancer cell line (PPC, LNCaP, DU145 and  
TSU) were assessed by Northern blot analysis using PCR radiolabeled probe. The result  
indicated that the transcript of the human counterpart was slightly smaller relative to the rat  
25 *p-Hyde* mRNA. The same blot was further hybridized with cyclophylin cDNA as internal  
control. In this autoradiogram, 28S and 18S ribosomal RNA were used as marker. The  
length of the transcript was calculated based on the Spirin's formula  $M = 1550 \times S^{2.1}$  where  
 $M$  = molecular weight and  $S$  = Svedberg's constant (McConkey, 1967). Signals of the *p*-  
*Hyde* and cyclophylin transcripts were quantitated and normalized using a computerized  
30 densitometer (Quantity-One software, PDI). The levels of *p-Hyde* mRNA expression from  
both rat and human prostate cancer cell lines were compared (Table I). The data indicated  
that there was differential expression of *p-Hyde* gene in both Dunning rat and human prostate  
cancer cell lines which suggests that there may be a functional correlation between *p-Hyde*  
expression and prostate cancer progression. MAT-Lu showed the highest level of  
35 transcription, whereas AT-3 had the lowest. These data also indicated that the level of

5 transcription in MAT-LyLu was relatively higher than that of AT-1, suggesting that the novel  
cDNA was the result of cDNA competition between MAT-LyLu and AT-1 cDNAs. More  
importantly, the human homologue of *p-Hyde* does exist as demonstrated in the human  
prostate cancer cell lines. The highest level of *p-Hyde* transcription occurred in PPC-1,  
whereas the lowest was in LNCaP cell line (Table 1).

10

**Analysis of the deduced amino acid sequence of *p-Hyde* derived from ORF:** Open  
reading of the cDNA consists of 1467 bases coding for 489 amino acid residues; the  
calculated molecular weight of this protein is 54.8 kD. Further molecular analyses of this  
deduced amino acid sequence is based on the Kyte-Doolittle hydrophilicity plot, James-Wolf  
15 antigenic index and Emini surface probability plot using Laser Gene DNASTar software. In  
addition, hydrophilicity profile of the deduced amino acid sequence was deduced based on  
Hopp and Woods using Antigen program for the prediction of its antigenic determinants  
(Table II). The results of the first three analyses were in agreement with the fourth one. Two  
peptide regions (residues 113 through 131 and residues 223 through 249) exhibited the  
20 highest points of hydrophilicity and the antigenic index. Both peptide sequences can be used  
as immunogenic peptide to generate antibody. Its application to detect the translation  
product of *p-Hyde* in Western analysis was in agreement with the highest score of its surface  
probability.

25

Table II.      Average Hydrophilicity      Amino-Acid Sequence

2.53	241 to 246
1.93	117 to 122
1.73	119 to 124

30

**Confirmation of the ORF by *in vitro* transcription and translation:** To confirm the reading  
frame based on the sequencing data, the *p-Hyde* cDNA which is constructed in pBluescript was  
digested with KpnI rendering the linearized construct to be accessible for T3-RNA polymerase.  
This enzyme directs the synthesis of the riboprobe with 5'cap structure (5'me7Gppp5'G analog)  
35 similar to those present in eukaryotic mRNA (Stratagene). The 5'cap is important to increase



5 the yield and the stability of the synthesized mRNA as well as to enhance the *in-vitro* translation efficiency of the mRNA. This capped mRNA was then used as a template for *in-vitro* translation using rabbit reticulocyte lysates (Stratagene) in the presence of <sup>35</sup>S-Methionine. The translation products were identified under denaturing conditions using polyacrylamide gel electrophoresis containing 8 M urea. The autoradiograph shown in Fig.13 revealed that two translation products  
10 (55 and 55.5 kDa protein) were observed.

Molecular weights of these two proteins are in agreement with the calculated molecular weight of the deduced amino acid sequence of the reading frame. In addition, two start codons were also identified in this reading frame correlating with the two translation products obtained  
15 through *in vitro* translation of the riboprobe. The difference in molecular weight between both translation products is in agreement with the difference in 4 amino acid residues (MSGE) on the C terminal between both start codons. The molecular weight difference between both translation products is in agreement with the 0.5 kDa molecular weight of MSGE.

20 **Subcloning of *p-Hyde* insert into pcDNA 3.1 (-) and transfection into rat and human prostate cancer cell lines:** Referring to the data of differential expression of *p-Hyde* in rat and human prostate cancer cell lines, it is tempting to integrate this gene into cell line expressing the lowest level of *p-Hyde* followed by the assessment of the function of this gene in its stable transfectants. For this purpose, *p-Hyde* cDNA was subcloned into pcDNA3.1 (-) mammalian  
25 expression vector (Invitrogen). The following features of pcDNA3.1 facilitate both subcloning and gene expression: (1) An extensive multiple cloning site to facilitate cloning in one direction. (2) A CMV promoter to drive the constitutive transcription and translation of the inserted *p-Hyde* cDNA, and (3) An SV40 splice acceptor linked to a SV40 polyadenylation signal to facilitate expression. The KpnI-XbaI fragment of the *p-Hyde* cDNA containing an intact open reading  
30 frame was successfully subcloned into pcDNA3.1 (-) vector. The CsCl-banding-purified-pcDNA containing HYDE insert (designated as pcHYDE) was then transfected into rat (AT-1 and AT-3) and human (DU145 and PPC-1) prostate cancer cell line using Lipofectamine (Gibco/BRL) under G418 selection. In addition to this pcHYDE, all cell lines were also  
35 successfully transfected with the pcDNA3.1(-) vector as negative control. The results of obtaining the stable transfectants indicated that pcHYDE gene product is not toxic to the host



5 cell.

10 ***In vitro* assessment of stable p- Hyde transfectants:** The function of pcHYDE in stable transfectants (AT1 and AT3 group) listed were assessed *in vitro* in association with the cell cycle and apoptosis. The objective for the apoptosis assessment is based on the hypothesis that *p-Hyde* is a rat homologue of the murine putative TSAP-6 gene which is associated with the upregulation of apoptosis response under the induction of tumor suppressor p53 (Amson et al., 1996). Whereas the cell cycle analysis is referred to the fact that MAT-LyLu cell line is highly metastatic and the level of *p-Hyde* expression in this cell line is relatively higher than AT-1 cell line.

15 **Cell cycle analysis:** The strategy of the cell cycle analysis is based on the arrest of cell population in G1 and S boundary after 24 hours treatment with 1 mM Hydroxyurea (Iwasaki et al., 1995). As a result, the G2 phase should be zero. Cells were harvested after hydroxyurea release at 0, 10 and 24 hours and subjected to flowcytometer analysis using standard propidium iodide staining. The result is shown in TABLE III and showed that both AT1-H1 and AT3-H1, at 10 hours after the release of Hydroxyurea, were relatively slow in entering the S-phase, whereas the parental cell line (AT1 and AT3) were faster. At 24 hours after the release, the G2 cells were all elevated indicating that cell cycle was in progress. Overall, the slow entrance into S phase in both pcHYDE stable transfectants does not corroborate directly that it is correlated with the slower tumor growth *in vivo*. The progress of re-entering the cell cycle at 0, 10 and 24 hours after the release of Hydroxyurea were followed by flowcytometer. Consistently, no G2 phase was detected at  $t_0$  - after 24 hours treatment with Hydroxyurea - demonstrating the cell cycle arrest in the G1 and S boundary.

30 Overall, the arrest of all cell cultures in G1 and S boundary by 1 mM Hydroxyurea treatment for 24 hours was confirmed referring to the 0% population of G2. These results demonstrate that the slower response in entering the S phase in AT1-H1 and AT3-H1 is likely due to the effect of the pcHYDE gene product that may reflect the growth characteristic of these stable transfectants *in vivo*.

35

5     **Induction of susceptibility to apoptosis:** Apoptotic response was assessed using DNA  
laddering assay. DNAs were extracted after the respective treatment (24 hours with 1 mM  
Hydroxyurea followed by 24 hours with 0.1 mM 5'-dFUrd) and analyzed on 1.6% agarose gel  
electrophoresis. In agreement with cell cycle analyses, apoptotic response of the stable  
transfectants AT1-H1 and AT3-H1 (pcHYDE transfected AT-1 and AT-3) are consistently and  
10     significantly higher relative to both parental (AT-1 and AT-3) and pcDNA-transfected parental  
cell lines (AT1-pc1 and AT3-pc1). In particular, the highest apoptotic response occurred in  
synchronized culture under the induction with 0.1 mM 5'-dFUrd.

15     The enhanced apoptotic response in AT1-H1 and AT3-H1 transfectant after hydroxyurea  
treatment is the result of "thymineless death" (Kyprianou, 1994, Kyprianou et al., 1994) leading  
to depletion of intracellular thymidine-5-triphosphate (TTP) pools through indirect inhibition of  
thymidylate synthetase by fluorodeoxyuridine. Taken together, these data demonstrate that the  
apoptotic response in the pcHYDE stable transfectants is due to the downstream effect of  
pcHYDE gene product.

20     **UV-damaged DNA cannot be repaired in prostate cells transfected with *p-Hyde*:** Three DNA  
enzyme repair systems were evaluated in parental compared to *p-Hyde* transfected cells: uridine  
phosphorylase, uridine kinase, and UV damage repair. UV damage repair was impaired in the  
*p-Hyde* transfected cells. Decreased DNA repair activity results in higher levels of intact  
25     photoproducts (64PP). Consistent with these data, *pHyde* transfected cells also had a significant  
reduction in survival following UV exposure compared to parental AT3 cells as determined by  
colony formation assay. Thus, DNA repair enzyme impairment correlated with shorter survival  
and induction of apoptosis in prostate cancer cells transfected with *p-Hyde*.

30     **Northern RNA analysis of *p-Hyde* confirms overexpression of *p-Hyde*:** To confirm the  
correlation between apoptotic response and *p-Hyde* on the transcription level, Northern analysis  
of the total RNA derived from AT3-H1, AT3-H2 relative to AT3 parental cell line was  
performed. The result clearly demonstrated that the transcription level of pcHYDE in stable  
transfectant of AT3-H1 and AT3-H2 were significantly higher relative to AT3 parental cell line.  
35     This Northern analysis was carried out only in AT3, AT3-H1 and AT3-H2 using Hyde PCR

5 product and internal control cyclophilin as a probe.

*p-Hyde* suppresses prostate cancer growth *in vivo*: The lowest level of *p-Hyde* expression was observed in AT3 cell line. For this reason, AT3 cell line was transfected with pcHYDE, a construct of *p-Hyde* in mammalian expression vector of pcDNA3.1(-) under G418 selection. As negative control, AT-3 cell line was also transfected with the vector only and the stable transfectants obtained was designated as AT3-pc.

15 The tumor growth of the parental cell line of AT-3 and in stable transfectant AT3-H1 and AT3-H2 have been evaluated *in vivo*. One million cells of each cell lines in 0.3 ml of Hanks solution were inoculated subcutaneously in each flank of inbred male Copenhagen rat. In this initial experiment, three groups of each five rats were injected with each cell line. The size of tumors were scored after a time schedule. These results indicated that both AT3-H1 and AT3-H2 stable transfectant grew significantly slower than the AT-3 parental cell line suggesting that the tumor growth regression in both stable transfectants are regulated by the pcHYDE gene product.

20 Tumor progression represents an accumulation of genetic changes that affect oncogene and tumor suppressor gene expression, thereby altering the responsiveness of the cell to autocrine and paracrine positive and negative growth regulators. The Dunning tumor rat model of prostate cancer tumor progression consists of a spectrum of prostate cancer phenotypes ranging from well differentiated to poorly differentiated with differing responses to androgens. Moreover, the sublines originated and evolved from the same original spontaneous rat prostate tumor, thus making this present study of tumor progression unique. The development of these cell lines as the results of progression of tumor within a single and multiple serial passage.

30 The screening strategy of competition hybridization of cDNA library generated from highly metastatic rat MAT-LyLu cell line resulted in a novel cDNA clone designated as *p-Hyde* (Rinaldy and Steiner, 1997). The full length of this cDNA, indicated as its restriction map, consists of 2713 nucleic acid residues; it contains two reading frames consisting of 1467 and 1452 residues, respectively. Forty six percent of the sequence is the untranslated region and majority of this is in the 3' end of the gene. The nucleic acid sequence of the rat *p-Hyde* is set

5        forth in SEQ ID NO 3 and the nucleic acid sequence of the human p-Hyde is set forth in SEQ ID NO 1.

10        The level of expression of *p-Hyde* was determined by Northern blot analyses in both rat and human prostate cancer cell lines and compared to the cyclophilin expression as internal control. The Dunning rat prostatic cancer cell line AT-1, MAT-LyLu, MAT-Lu, AT-3 and G and human prostate cancer cell lines PPC1, LNCaP, TSU and DU145 are all positive with different level of expression (Table I). The transcript size of the human counterpart is slightly smaller relative to the rat *p-Hyde* mRNA. The data also indicated that there is a differential expression of *p-Hyde* gene in both rat and human prostatic cancer cell lines suggesting that there may be a functional correlation between *p-Hyde* expression and prostate cancer progression. MAT-Lu showed the highest level of transcription, whereas AT-3 had the lowest. The striking difference between these two cell lines is important in its association with the growth characteristic of MAT-Lu cell line which is highly metastatic. These data also indicated that the level of transcription in MAT-LyLu was relatively (10%) higher than that of AT-1, demonstrating that the novel cDNA was the result of cDNA competition between MAT-LyLu and AT-1 cDNAs (Rinaldy and Steiner, 1997). Of significance, the human homologue of *p-Hyde* does exist in the human prostate cancer cell lines. The highest level of *p-Hyde* transcription occurred in PPC1, a primary androgen sensitive human prostate cancer cell line, whereas it was lowest in LNCaP cell line, an androgen-insensitive prostate cancer cell line. Thus, *p-Hyde* does not appear to be specific to mammalian prostate tissue, but has been found in other tissues including placenta and breast and in other species such as mouse and human suggesting that its role is important in fundamental cellular biology.

30        **Tumor suppression and induction of susceptibility to apoptosis:** Interestingly, *p-Hyde* has the dual ability to act like a tumor suppressor gene and induce susceptibility to apoptosis by what may be p53 independent pathways. The growth of prostate tumors in rats was greatly inhibited by *p-Hyde*. Moreover, prostate cancer cells expressing *p-Hyde* were more sensitive to UV DNA damage driving these cells into cell programmed death. Analysis of DNA repair enzyme activity suggests a defect resulting in the presence of intact (6-4) PP and decreased cell survival by colony forming assay. However, the capacity of *p-Hyde* to induce susceptibility to apoptosis

5 is not limited to UV DNA damage. Chemotherapy agent, Fluorodeoxyuridine, a pyrimidine  
antimetabolite which is related to fluorouracil (5-FU) and has been used for treatment of a wide  
variety of human epithelial malignancies, also more readily induces apoptosis in prostate cancer  
cell expressing *p-Hyde*. Moreover, cancer cells expressing *p-Hyde* are also more susceptible to  
10 gamma radiation. Thus, the mechanisms of cellular DNA injury are different for UV, gamma  
radiation, and Fluorodeoxyuridine suggesting that the ability to make cells more susceptible to  
apoptosis is more global in action. This unique function of *p-Hyde* represents a new class genes  
that induce susceptibility to apoptosis. This is different than the function ascribed to tumor  
suppressor genes like p53 which directly induces apoptosis, not sensitivity to apoptosis (Yonish-  
15 Rouach *et al.*, 1991). Moreover, *p-Hyde* activity is in contrast to *bcl-2* where the absence of *bcl*-  
2, not the presence of, makes the cancer cell more susceptible to cell programmed death  
(McDonnellet *et al.*, 1992).

Use in therapy of human disease: These unique functional features of the *p-Hyde* gene may  
be exploited for the treatment of hyperproliferative disorders and cancer. One effective  
20 therapeutic strategy, for example, may be the treatment of carcinoma cells expressing *p-Hyde*  
with chemotherapy agents or UV mimetic drugs (such as acetylaminofluorine). However, cancer  
cells are not likely to produce significant levels of the growth inhibition *p-Hyde*. Consequently,  
the *p-Hyde* gene may be introduced into cancer cells by gene therapy. Tumors transduced with  
vectors containing *p-Hyde* may not only be directly suppressed by *p-Hyde* as demonstrated in  
25 this study, but also when treated in combination with DNA damaging therapy such as  
chemotherapy, UV mimetic drugs, or radiation, have even a greater anti-cancer effect. Since gene  
therapy will target cancer cells, then enhancement of apoptosis will occur more selectively in  
cancer cells following DNA damage (UV, radiation, or chemotherapy).

30  
**EXAMPLE 2: Prostate Cancer Gene Therapy Using Adenovirus Expressing A Novel  
Tumor Suppressor Gene pHyde**



## MATERIALS AND METHODS

**Cell lines and tissue culture condition:** Human prostate cancer cell lines (obtained from ATCC, Rockville, Maryland) PPC-1, DU145, PC-3, LNCaP, and TSU-Pr1 were grown in RPMI-1640 medium (Cellgro, Herndon, VA) containing 10% fetal bovine serum (Hyclone Laboratories, Logan, UT) at 37°C and 5% CO<sub>2</sub>. Human embryonic kidney cell line 293 (ATCC) was grown in D-MEM medium (Cellgro) containing 10% heat inactivated fetal bovine serum at 37°C and 5% CO<sub>2</sub>.

**Construction of AdRSVpHyde:** A rat pHyde cDNA gene was isolated as described in U.S Serial No: 09/302,457. After digestion with EcoR I, a 2.6 kb fragment which contains the 1467 bp full-length coding sequence of pHyde cDNA was subcloned under the control of a truncated RSV promoter (395 bp) into an E1/E3 deleted adenoviral shuttle vector. The resultant adenoviral shuttle vector was cotransfected into 293 cells with pJM17, an adenoviral type 5 genome plasmid, by calcium phosphate method. Individual plaques were screened for recombinant AdRSVpHyde by PCR using specific primers for both the RSV promoter and pHyde cDNA sequences. Single viral clones were propagated in 293 cells. The culture medium of the 293 cells showing the completed cytopathic effect (CPE) was collected, and the adenovirus was purified and concentrated by twice CsCl<sub>2</sub> gradient ultracentrifugation. The viral titration and transduction were performed as previously described. The schematic diagram of AdRSVpHyde was illustrated in Fig. 1. The sequence of AdRSVpHyde is set forth in Figure 10.

**Northern blot:** Cells were extracted and total RNA was isolated by RNeasy Kit (Qiagen, Santa Clarita, CA). Total RNA was loaded on a 1.2% polyacrylamide gel and processed to electrophoresis. The standard Northern blot transfer to a Nylon membrane (Hybond-N<sup>+</sup>, Amersham Life Science, Buckinghamshire, England) was performed as previously described. The cDNA probes (pHyde or p53) were labeled by α-<sup>32</sup>P-dCTP using random primer method (Prime-It II Kit, Stratagene, La Jolla, CA). The membrane was hybridized with the probe in Rapid-hyb buffer (Amersham Life Science) according to the Manufacturer's protocol. The membrane was exposed to a Kodak X-ray film under one intensifying screens at -80°C for autoradiography. GAPDH cDNA probe was labeled as described above and used as an internal control for normalization.

5      **Western blot:** Cells were extracted and processed for gel electrophoresis as previously  
described.( Lu Y, Whitaker L, Li X, et al. Coexpression of galectin-1 and its complementary  
glycoconjugates laminin and lysosome-associated membrane proteins in murine PCC4.aza1R  
embryonal carcinoma cells induced to differentiation by butyrate. *Mol Cell Differ.* 1995;3:175-  
191). Cell extract lysates (100 mg) were loaded on 12% polyacrylamide gels and subjected to  
10      sodium dodecylsulfate (SDS) gel electrophoresis, then transferred to a nitrocellulose membrane  
(Bio-Rad Laboratories, Hercules, CA). The membrane was treated with blocking solution (15%  
nonfat milk, 0.02% sodium azide in phosphate-buffered saline) overnight at 4°C. The membrane  
was incubated for 1 hr at room temperature with rabbit anti-rat pHyde polyclonal antibody  
(specifically generated by Research Genetics, Inc. based on the computer-created antigenic  
15      peptide derived from pHyde coding sequences). The membrane was then incubated for 1 hr at  
room temperature with <sup>125</sup>I-labeled second antibody (Amersham Life Science, Arlington Heights,  
IL). The membrane was exposed to a Kodak X-ray film between two intensifying screens at -  
80°C for autoradiography.

20      **AdRSVpHyde in vitro studies:** Human prostate cancer cells were infected with AdRSVpHyde  
*in vitro* with a multiplicity of infection (MOI) of 100 or 200. After viral infection, cells were  
incubated at 37°C and cell numbers were determined at day 5 after viral infection.

25      **In vivo studies using DU145 xenograft tumors:** DU145 cells ( $1.4 \times 10^7$  cells in 0.2 ml of PBS)  
were injected subcutaneously into the flank of male nude mice (Harlan Sprague Dawley). When  
the tumors reached an average volume of 80 mm<sup>3</sup>,  $5 \times 10^9$  pfu adenoviral vectors (AdRSVpHyde  
or control adenovirus AdRSVlacZ) or PBS alone for untreated controls were injected directly  
into the tumor. Tumor volume was measured every three to four days until the animals were  
euthanized. All the animals were sacrificed at day 52 after viral injection when several of them  
30      showed distress or had a tumor burden greater than 15% of total body weight. Tumor samples  
were collected and processed for H&E staining.

5     **DNA extraction and gel electrophoretic analysis of DNA fragmentation:** Soluable DNA was  
extracted as described previously (in Oridate N, Lotan D, Xu X-C, Hong WK, and Lotan R.  
Differentiation induction of apoptosis by all-trans-retinoic acid and N-(4-  
hydroxyphenyl)retinamide in human head and neck squamous cell carcinoma cell lines. *Clin.*  
10     *Cancer Res.* 1996;2:855-863) Briefly, the cells floating in medium were collected 48 h post  
transduction by centrifugation. The pellet was resuspended in Tris-EDTA buffer (pH 8.0). The  
cells were lysed in 10 mM Tr-s-HCl (pH 8.0), 10 mM EDTA, and 0.5% Triton X-100 on ice for  
15 min. The lysate was centrifuged at 12,000xg for 15 min to separate soluble (fragmented)  
DNA from pellet (intact genomic) DNA. Soluble DNA was treated with Rnase A (50 ug/ml) at  
37C for 1 h, followed by treatment with proteinase K (100 ug/ml) in 0.5% SDS, at 50C for 2 h.  
15     The residual material was extracted with phenol/chloroform, precipitated in ethanol,  
electrophoresed on a 2% agarose gel.

## RESULTS

20     **Exogenous pHyde expression in DU145 cells:** To determine whether AdRSVpHyde is able to  
successfully transfer and express rat pHyde at mRNA and protein levels, DU145 cells were  
transduced by AdRSVpHyde at MOI=200. The cell extract were harvested 48 h after viral  
transduction to determine the pHyde expression. While there was a minor endogenous  
expression of pHyde at mRNA level (Fig. 2A) but not at protein level (Fig. 2B) in DU145 cells,  
25     there was an apparent high exogenous pHyde expression induced by AdRSVpHyde at both  
mRNA (Fig. 2A) and protein (Fig. 2B) levels.

30     **Prostate cancer cell growth inhibited by AdRSVHyde:** To determine the effects of pHyde on  
cell growth of human prostate cancer cell lines, DU145 and LNCaP were treated with  
AdRSVpHyde, AdRSVlacZ (control vector), and no virus *in vitro*. AdRSVpHyde significantly  
inhibited the growth of DU145 and LNCaP cells, with 76.9% (Fig. 3A) and 83.1% (Fig. 3B)  
inhibition respectively, compared to untreated control cells.

35     **AdRSVpHyde inhibited prostate tumor growth in vivo:** To evaluate the effects of AdRSVpHyde  
treatment of prostate cancer cell growth *in vivo*, DU145 human prostate tumors were established

5 in nude mice by injecting  $1.4 \times 10^7$  PPC-1 cells subcutaneously into the flanks of nude mice. When mice developed tumors with an averaging  $80 \text{ mm}^3$  volume, the mice were divided into three groups: AdRSVpHyde treated ( $n=7$ ), AdRSVlacZ control virus treated ( $n=7$ ), and untreated groups ( $n=7$ ). Treated tumors were injected with a single dose of  $5 \times 10^9$  pfu of either the control virus or AdRSVpHyde. As shown in Fig. 4, untreated and control virus treated DU145 tumors  
10 grew rapidly relative to the AdRSVpHyde treated tumors. By day 53 following viral injection, the tumor burden in nude mice bearing untreated and control virus treated DU145 tumors reached  $5953 \text{ mm}^3$  and  $4777 \text{ mm}^3$  respectively. In contrast, DU145 tumors transduced by AdRSVpHyde had a significant reduction in tumor volume ( $1515 \text{ mm}^3$ ) compared to untreated

5 and control virus treated-DU145 tumors, that is, 25.4% of untreated and 31.7% of control virus treated DU145 tumor volume (Fig. 4).

*Growth inhibition by AdRSVpHyde correlated to p53 expression in prostate cancer cells:* One observation in the *in vitro* study was that the phenotype of DU145 and LNCaP cells transduced by AdRSVpHyde were also altered. Unlike the control and control virus treated cells, AdRSVpHyde-transduced cells had a marked morphology of round, detaching and floated dying cells, a characteristic of cells undergoing apoptosis, and cell number decreased quickly over time (Fig. 5). Consistently, AdRSVpHyde showed a strong inhibition on the growth of these two cell lines (Fig. 3). However, AdRSVpHyde did not inhibit the cellular growth of PC-3 and TSU-Pr cells, and had only a minor growth inhibition on PPC-1 cells (data not shown). Consistently, there was no evident morphological differences between untreated control, control virus-treated and AdRSVpHyde-treated cells in these lines (Fig. 6). To determine whether the differential expression of various genes, especially those involved in apoptosis pathway, accounted for the differential inhibitory effect by AdRSVpHyde on different prostate cancer cell lines, several genes including p53 and Rb were screened at mRNA level by Northern hybridization. Interestingly but not too surprisingly, p53 was found to only express in DU145 and LNCaP cells but not in PC-3, TSU-Pr, and PPC-1 cells; in contrast, Rb gene were all expressed at the mRNA level in these cells (Fig. 7). Therefore, it appeared that there was a correlation between p53 expression and AdRSVpHyde-mediated inhibition. To determine whether pHyde regulated p53 expression, the same Northern blot in Fig. 2A was stripped and rehybridized with p53 probe. Indeed an induction of p53 mRNA was observed in AdRSVpHyde transduced DU145 cells (Fig. 8). Furthermore, transduction of LNCaP cells by AdRSVpHyde showed DNA laddering pattern, a marker for cells undergoing apoptosis (Fig. 9), indicate that pHyde expression induced apoptosis in LNCaP cells. Taken together, these results demonstrate that pHyde may function via p53 pathway to induce apoptosis and its growth inhibition may depend on p53 expression in the target cells.

The previous study showed that pHyde has the dual ability to act like a tumor suppressor gene and induce susceptibility to apoptosis. The growth of prostate tumors in rats was greatly inhibited by p-Hyde. Moreover, prostate cancer cells expressing pHyde were more sensitive to UV DNA



5 damage driving these cells into cell programmed death. In this study AdRSVpHyde was shown  
to have an effective inhibition on cell growth both *in vitro* and *in vivo* for the prostate cancer  
cells expressing p53, but not for the prostate cancer cells missing p53 expression. One  
possibility is that pHyde can induces apoptosis by p53 dependent pathway (such as in DU145  
and LNCaP cells) and p53 independent pathway which requires an outside cell death trigger such  
10 as UV or chemical (such as methylnitrosourea) to act as a co-inducer for apoptosis. Consistent  
with our result that only DU145 and LNCaP, but not PPC-1, PC-3 and TSU-Pr, expressed p53  
at mRNA level (Fig. 10), other groups found that only DU145 and LNCaP cells expressed p53  
at protein level but not PC-3 and TSU-Pr. Interestingly, p53 protein in DU145 was claimed to  
be a mutant p53.<sup>21</sup> Therefore, it seems that existence of p53 protein, regardless of its wild-type  
15 or mutant status, is required for pHyde to act as a tumor suppressor gene alone. The possibility,  
that the mutation in p53 protein in DU145 cells may not affect its ability for pHyde-mediated  
growth inhibition, need to be further studied. Moreover, the potential inhibitory effects by  
combination of AdRSVpHyde and UV (or methylnitrosourea) on cells missing p53 protein  
expression (such as PC-3 and TSU-Pr) will be further characterized.

20 One interesting and important finding of this study was that AdRSVpHyde induced p53  
expression in DU145 cells. Not only this may explain that pHyde acts in a more global manner  
to induce susceptibility to apoptosis, but also partially explains why pHyde has a partial sequence  
homologue with TSAP-6, a human protein claimed to be involved in p53-associated pathway:  
25 because pHyde could be a rat homologue of TSAP-6 gene, or a member of the TSAP-6 like  
family which is involved in p53-associated pathway. Furthermore, there is very few, if any,  
identified cellular proteins which act as regulators for p53 gene, which is a transcription factor  
to act many downstream genes. The significance of the finding that pHyde up-regulates p53  
expression and the consequent exploration for a new cellular regulation mechanism are exciting.  
30 Nevertheless, whether pHyde directly regulates p53 gene at the transcriptional level will be  
determined in our current study by employing p53 promoter/CAT reporter chimeric gene in the  
presence and absence of pHyde protein in DU145 cells.

In summary, pHyde is a novel a tumor suppressor gene and AdRSVpHyde effectively inhibites  
prostate cancer both *in vitro* and *in vivo*. The monotherapy of AdRSVpHyde alone or combined  
35 therapy of AdRSVpHyde with radio- or chemo-therapy should have an effective therapeutic

potential for treatment of locally advanced prostate cancer.